

PSB NASA  
IN- 91-CR  
67682  
P 55

## Shock Wave Data for Minerals

Thomas J. Ahrens and Mary L. Johnson

### 1. INTRODUCTION

Shock compression of the materials of planetary interiors yields data, which upon comparison with density-pressure and density-sound velocity profiles of both terrestrial planetary mantles and cores [4,5,94], as well as density profiles for the interior of the major planets [148], constrain internal composition and temperature. Other important applications of shock wave data and related properties are found in the impact mechanics of terrestrial planets and the solid satellites of the terrestrial and major planets. Significant processes which can, or have been, studied using shock wave data include: (1) the formation of planetary metallic cores during accretion [169,192], and (2) the production of a shock-melted "magma ocean" and concurrent impact volatilization versus retention of volatiles during accretion [1]. Also of interest are the shock-induced chemical reactions between meteoritic components (e.g. H<sub>2</sub>O and Fe: [111]). The formation of primitive atmospheres, for example, containing a large fraction of H<sub>2</sub>O and CO<sub>2</sub> is also addressable using shock wave and other thermodynamic data for volatile-bearing minerals (e.g. [110,112]). A related application of both shock

(NASA-CR-199490) SHOCK WAVES DATA  
FOR MINERALS (California Inst. of  
Tech.) 55 p

N96-133350  
Unclassified

63/91 0067682

May 9, 1994

compression and isentropic release data for minerals [13,14] is in the mechanics of both the continued bombardment and hence cratering on planetary objects through geologic time [170], as well as the effects of giant impacts on the Earth [183,185]. Finally, recovery and characterization of shock-compressed materials have provided important insights into the nature of shock deformation mechanisms and, in some cases, provided physical data on the nature of either shock-induced phase changes or phase changes which occur upon isentropic release from the high-pressure shock state (e.g., melting) [193,194].

As indicated for the data summary of Table 1, a very large data set exists describing the Hugoniot equation of state of minerals. Whereas some earlier summaries have provided raw shock data [47,121,213], the present summary provides fits to shock wave data. Earlier summaries providing fits to data are given by Al'tshuler et al. [24] and Trunin [203].

Hugoniot data specify the locus of pressure-density (or specific volume) states which can be achieved by a mineral from some initial state with a specified initial density. An analogous summary for rocks, usually described as a mixture of minerals are given in Chapter 3-4.

Three pressure units are commonly in use in the shock wave literature: kilobar (kbar), gigapascal (GPa), and megabar (Mbar). These are equal to  $10^9$ ,  $10^{10}$ , and  $10^{12}$  dyne/cm<sup>2</sup>, respectively, or  $10^8$ ,  $10^9$ , and  $10^{11}$  pascals in SI units.

## 2. SHOCK WAVE EQUATION OF STATE

The propagation of a shock wave from a detonating explosive or the shock wave induced upon impact of a flyer plate accelerated, via explosives or with a gun, result in nearly steady waves in materials. For steady waves a shock velocity  $U_s$  with respect to the laboratory frame can

be defined. Conservation of mass, momentum, and energy across a shock front can then be expressed as

$$\rho_1 = \rho_0 (U_s - u_0) / (U_s - u_1) \quad (1)$$

$$P_1 - P_0 = \rho_0 (u_1 - u_0) (U_s - u_0) \quad (2)$$

$$E_1 - E_0 = (P_1 + P_0)(1/\rho_0 - 1/\rho_1)/2 = 1/2 (u_1 - u_0)^2 \quad (3)$$

where  $\rho$ ,  $u$ ,  $P$ , and  $E$  are density, particle velocity, shock pressure, and internal energy per unit mass and, as indicated in Fig. 1, the subscripts 0 and 1 refer to the state in front of and behind the shock front, respectively. In Table 1, shock velocity and particle velocity are designated as  $U_s$  and  $U_p$ . Equations (1)-(3) are often called the Rankine-Hugoniot equations. It should be understood that in this section pressure is used in place of stress in the indicated wave propagation direction. In actuality, stress in the wave propagation direction is specified by Eq. (2). A detailed derivation of Eqs. (1), (2), and (3) is given in Duvall and Fowles [70]. Equation (3) also indicates that the material achieves an increase in internal energy (per unit mass) which is exactly equal to the kinetic energy per unit mass.

In the simplest case, when a single shock state is achieved via a shock front, the Rankine-Hugoniot equations involve six variables ( $U_s$ ,  $u_1$ ,  $\rho_0$ ,  $\rho_1$ ,  $E_1 - E_0$ , and  $P_1$ ); thus, measuring three, usually  $U_s$ ,  $u_1$ , and  $\rho_0$ , determines the shock state variables  $\rho_1$ ,  $E_1 - E_0$  and  $P_1$ . The key assumption underpinning the validity of Eqs. (1)-(3) is that the shock wave is steady, so that the rise time  $\tau_s$  is short compared to the characteristic time for which the high pressure, density, etc. are constant (see Fig. 1). Upon driving a shock of pressure  $P_1$  into a material, a final shock state is achieved which is described by Eqs. (1)-(3). This shock state is shown in Fig. 2, in relation to other thermodynamic paths, in the pressure-volume plane. Here  $V_0 = 1/\rho_0$  and  $V = 1/\rho$ . In the case of the isotherm and isentrope, it is possible to follow, as a thermodynamic path, the actual isothermal or isentropic curve to achieve a

state on the isotherm or isentrope. A shock, or Hugoniot, state is different, however. The Hugoniot state ( $P_1, V_1$ ) is achieved via a shock front. The initial and final states are connected by a straight line called a Rayleigh line (Fig. 2). Thus successive states along the Hugoniot curve cannot be achieved, one from another, by a shock process. The Hugoniot curve itself then just represents the locus of final shock states corresponding to a given initial state.

It has long been recognized that the kinematic parameters measured in shock wave experiments  $U_s$  and  $U_p$  can empirically be described in regions where a substantial phase change in the material does not occur as:

$$U_s = C_0 + S U_p \quad (4)$$

As further discussed in several review articles on shock compression [22,59,136], and a recent book [40], Hugoniot data for minerals and other condensed media may be described over varying ranges of pressure and density in terms of a linear relation of shock and particle velocity in Table 1. This table was assembled using the *Microsoft Excel*, version 3.0, (Redmond, WA 1993) program and the least-square fits to the shock wave data with standard errors were derived by using the LINEST function. The equations employed for line slopes and intercepts are identical to those given in Bevington [46] (Eq. 6-9, p. 104; Eq. 6-21, p. 114 for errors in slopes; and Eq. 6-22, p. 114 for errors in intercepts).

The  $U_s$ - $U_p$  data for a wide range of minerals are given in Table 1. Here  $C_0$  is the shock velocity at infinitesimally small particle velocity, or the ambient pressure bulk sound velocity which is given by

$$C_0 = \sqrt{K_s/\rho_0}, \quad (5)$$

where  $K_s$  is the isentropic bulk modulus. (Bevington [46]

$K_S = -V \frac{dP}{dV}_S$  in the absence of strength effect (see Sect. 3). Upon substituting Eq. 4 into Eq. 2, and denoting the shock pressure as  $P_H$ , this is given by:

$$P_H = \rho_0 U_p (C_0 + S U_p) \quad (6)$$

Thus, from the form of Eq. 6, shock pressure is given as the sum of a linear and quadratic term in particle velocity, based on the data of Table 1. A pressure-volume relation can be obtained by combining Eq. 6 with Eq. 1 to yield:

$$P_H = \rho_0 C_0^2 \eta / (1 - S\eta)^2 \quad (7)$$

where

$$\eta = 1 - V/V_0 = U_p / U_S. \quad (8)$$

Eq. 7 is often called the "shock wave equation of state" since it defines a curve in the pressure-volume plane.

The isentropic pressure can be written (e.g. [93,98,171]) by an expression analogous to Eq. 7 as a series

$$P_S = \rho_0 C_0^2 (\eta + 2S\eta^2 + \dots), \quad (9)$$

which upon differentiation yield the isentropic bulk modulus

$$K_S = \rho_0 C_0^2 (1 + (4S-1)\eta + \dots) \quad (10)$$

The analogous bulk modulus along the Hugoniot is:

$$K_H = -V \frac{\partial P}{\partial V} |_H. \quad (11)$$

The isentrope and the Hugoniot and isentropic bulk modulus are related via:

$$K_S = K_H + \left(\frac{\gamma}{2}\right) [P_H - K_H \eta / (1-\eta)] - [P_H - P_S] [\gamma + 1 - q_0 (1 - q'\eta + \dots)] \quad (12)$$

Here we assume a volume dependence of the Gruneisen parameter

$$\gamma = V (\partial P / \partial E)_V = \gamma_0 (V/V_0)^q, \quad (13)$$

where

$$q = d \ln \gamma / d \ln V \text{ and } q' = d \ln q / d \ln V \quad (14)$$

$\gamma_0$  is the Gruneisen parameter under standard pressure and temperatures and is given by

$$\gamma_0 = \alpha K_T V_0 / C_V = \alpha K_S V_0 / C_P, \quad (15)$$

where  $\alpha$  is the thermal expansion coefficient,  $K_T$  is the isothermal bulk modulus and  $C_P$  and  $C_V$  are the specific heat at constant pressure and volume. We note that the  $P_S$  and  $P_H$  can be related by assuming the Mie-Gruneisen relation

$$P_H - P_S = \frac{\gamma}{V} (E_H - E_S), \quad (16)$$

if  $\gamma$  is independent of temperature, where  $E_H = E_1 - E_0$  is given by Eq. 3 and  $E_S$  is given by

$$E_S = - \int_{V_0}^V P_S dV. \quad (17)$$

Because the Gruneisen ratio relates the isentropic pressure,  $P_S$ , and bulk modulus,  $K_S$ , to the Hugoniot pressure,  $P_H$ , and Hugoniot bulk modulus,  $K_H$ , it is a key equation of state parameter.

The shock-velocity particle relation of Table 1 can be used to calculate the shock pressure when two objects impact. If (A) the flyer plate and (B) the target are known and expressed in the form of Eq. (7), the particle velocity  $u_1$  and pressure  $P_1$  of the shock state produced upon impact of a flyer plate at velocity  $u_{fp}$  on a stationary target may be calculated from the solution of the equation equating the shock pressures in the flyer and driver plate:

$$\frac{\rho_{oA} (u_{fp} - u_1)(C_{oA} + S_A (u_{fp} - u_1))}{\rho_{oB} u_1 (C_{oB} + S_B u_1)} = 1 \quad (18)$$

That is;

$$u_1 = \frac{(-b - \sqrt{b^2 - 4ac})}{2a}, \quad (19)$$

where

$$a = S_A \rho_{oA} - \rho_{oB} S_B, \quad (20)$$

$$b = C_{oA} \rho_{oA} - 2S_A \rho_{oA} u_{fp} - \rho_{oB} C_{oB}, \quad (21)$$

and

$$c = u_{fp} (C_{oA} \rho_{oA} + S_A \rho_{oA} u_{fp}). \quad (22)$$

### **3. SHOCK-INDUCED DYNAMIC YIELDING AND PHASE TRANSITIONS**

Both dynamic yielding and phase transitions give rise to multiple shock wave profiles when pressure or particle velocity versus time is recorded. Virtually all nonporous minerals and rocks in which dynamic compression has been studied demonstrate phenomenon related to dynamic yielding, in which materials transform from finite elastic strain states to states in which irreversible deformation has occurred. Moreover, most minerals and a large number of

compounds, elements, and organic materials demonstrate shock-induced phase changes.

The dynamic yield point under shock compression, the Hugoniot elastic limit, or HEL, is defined as the maximum shock pressure a material may be subjected to without permanent, massive, microscopic rearrangement taking place at the shock front. As shown in Fig. 3a, the shock velocity of the HEL state remains nearly constant and for non-porous media is usually equal to the longitudinal elastic wave velocity. Viscoelastic polymeric media generally do not display the HEL phenomenon. We denote five regimes in Fig. 3 for the case of dynamic yielding and phase transition and the available shock wave data are separately fit to linear relations in these regimes in Table 1. For some minerals there are more than four regimes indicated, for reasons such as crystallographic control of compression at low pressures (such as O<sub>a</sub>, O<sub>b</sub> for quartz), and for more than one high-pressure state (such as 4a, 4b, and 4c for halite).

The crystallographic or atomistic level nature of shock-induced phase changes varies from simple average coordination changes observed in various liquids, ionization and debonding in non-metallic fluids, electronic transitions in metal and non-metals, changes in crystal structure in solid materials, and transition from the solid to the fluid state.

In the case of a phase change, the pressure along the isentrope P<sub>s</sub> at the volume V<sub>1</sub> corresponding to a Hugoniot state (P<sub>1</sub>, V<sub>1</sub>) is given by

$$\frac{P_1}{2} (V_{\infty} - V_1) = - \int_{V_0}^{V_1} P dV + \frac{V_1}{\gamma} (P_1 - P_s) + ETR \quad (23)$$

where the left-hand side is the Rankine-Hugoniot energy, and the first and second terms on the right represent the gain in the internal energy along the paths 1 and 2 of Fig. 2. Here V<sub>∞</sub> is the specific volume of the initial material and V<sub>0</sub> the specific volume of the shock-induced high-pressure phase, or the intrinsic volume of the sample if the

initial state is distended. Also  $E_{TR}$  is the energy of transition to the high-pressure phase at STP. In the case of no phase change,  $E_{TR} = 0$ . For zero initial porosity  $V_{00} = V_0$ . The unknown parameter in Eq. 23 is  $P_s$ , which is implicit in the first integral term on the right-hand side and explicit in the second term. The second term is obtained by using the definition of the Gruneisen parameter (Eq. 13) to calculate the change in energy associated with the pressure difference ( $P_1 - P_s$ ) at constant volume.

#### 4. SHOCK TEMPERATURES

For many condensed media, the Mie-Gruneisen equation of state, based on a finite-difference formulation of the Gruneisen parameter (Eq. 16), can be used to describe shock and postshock temperatures. The temperature along the isentrope [224] is given by

$$T_s = T_i \exp \left[ - \int \frac{V_b}{V_a} \left( \frac{\gamma}{V} \right) dV \right] \quad (24)$$

where  $T_i$  is the initial temperature. For the principal isentrope centered at room temperature,  $T_i = T_0$ ,  $V_a = V_0$ , initial volume, and  $V_b = V$ , compressed volume. For the calculation of postshock temperatures  $T_i = T_H$ , the Hugoniot temperature,  $V_a = V_H$ , the volume of the shock state, and  $V_b = V_{00}$ , the postshock volume corresponding to the postshock temperature. For shock compression to a volume  $V$ ,  $P_s$  is first obtained by using Eq. 23; then  $T_s$ , the isentropic compression temperature at volume  $V$ , may be calculated by using Eq. 24. Finally, using Eq. 16, the shock temperature  $T_H$  is given by

$$\frac{V}{\gamma} (P_H - P_s) = \int_{T_s}^{T_H} C_V dT \quad (25)$$

It is useful to carry out both postshock and shock temperature measurements as they provide complementary information for the thermal equation of state, that is,  $\gamma$ , as well as  $C_V$ . Minerals for which shock temperatures have been (usually via radiative techniques) are so indicated in Table 1.

In the case of molecular fluids such as water, a formulation based on the near constancy of  $C_P$  at constant pressure is used [41,167].

Although there have been few data collected, postshock temperatures are very sensitive to the models which specify  $\gamma$  and its volume dependence, in the case of the Gruneisen equation of state [49,164,165]. In contrast, the absolute values of shock temperatures are sensitive to the phase transition energy  $E_{TR}$  of Eq. 23, whereas the slope of the  $T_H$  vs. pressure curve is sensitive to the specific heat.

*Acknowledgments.* Research supported under NSF, NASA, and DoD. We appreciate comments on this manuscript from William W. Anderson, Kathleen Gallagher, Wenbo Yang, and J. Michael Brown. Contribution number 5332, Division of Geological and Planetary Sciences.


#### **FIGURE CAPTIONS**

FIG. 1. Profile of a steady shock wave, rise time  $\tau_s$ , imparting a particle velocity  $u_1$  pressure  $P_1$ , and internal energy density  $E_1$ , propagating with velocity  $U_s$  into material that is at rest at density  $\rho_0$  and internal energy density  $E_0$ .


FIG. 2. Pressure-volume compression curves. For isentrope and isotherm, the thermodynamic path coincides with the locus of states, whereas for shock, the thermodynamic path is a straight line to point  $P_1, V_1$ , on the Hugoniot curve, which is the locus of shock states.

FIG. 3. Sketch of shock velocity-particle relation (a) and corresponding pressure-volume Hugoniot curves (b) for a mineral which undergoes dynamic yielding and a phase change.

- 0: compression up to the Hugoniot Elastic Limit (HEL)
- 1: transition via dynamic yielding to a quasi-hydrostatic state
- 2: low pressure state
- 3: mixed region
- 4: high pressure state



TJAS4153SFD



TJA84154SFD




TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
<b>Gases:</b>											
Air <sup>c</sup>	(mixture)	0.884	2.28	--	1.20	--	4.317	5.788	2	2	[147]
			4.25	--	0.85	--	5.788	7.379	4	2	
Nitrogen plus Oxygen <sup>c</sup>	1:1 N <sub>2</sub> +O <sub>2</sub>	0.945	1.83	0.11	1.26	0.03	2.235	3.785	2	6	[179]
Nitric Oxide <sup>e</sup>	NO	1.263	3.76	0.06	0.98	0.02	2.01	3.245	2	7	[179]
Ammonia <sup>i</sup>	NH <sub>3</sub>	0.715	2.45	0.19	1.34	0.04	1.01	7.566	2	12	[83,121,140]/[159]
Argon <sup>k</sup>	Ar	0.0013	0.71	0.10	1.041	0.018	1.73	7.81	2	25	[58,71,84]
Argon <sup>f</sup>	Ar	0.919	1.04	0.06	1.36	0.02	1.59	4.04	2	6	[214]
Argon <sup>l</sup>	Ar	1.026	1.1	0.2	1.45	0.07	1.42	4.10	2	7	[213]
Argon <sup>c</sup>	Ar	1.401	1.01	0.10	1.79	0.08	0.301	1.35	2	9	
			1.28	0.06	1.58	0.02	1.32	3.758	3	24	
			3.04	0.14	1.09	0.03	3.65	6.451	4	10	[80,121,146,180, 214] / [80,221]
Argon <sup>c</sup>	Ar	1.65	0.88	0.15	2.00	0.11	0.56	1.85	2	8	
			1.9	0.3	1.46	0.10	1.85	3.60	3	7	
			2.5	0.3	1.29	0.07	3.60	4.60	4	3	[64,109,121,189]
Carbon Dioxide <sup>i</sup>	CO <sub>2</sub>	1.173	1.54	0.09	1.44	0.03	1.585	3.765	2	16	
			3.3	0.4	1.01	0.07	4.549	6.264	4	3	[147,178]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Carbon Dioxide <sup>h</sup>	CO <sub>2</sub>	1.541	1.99 3.27	0.08 --	1.56 1.21	0.03 --	1.03 3.68	3.68 4.79	2 4	3 2	[233]
Carbon Monoxide <sup>c</sup>	CO	0.807	1.54 2.59 1.3	-- 0.04 0.3	1.40 0.974 1.21	-- 0.010 0.04	1.692 2.471 5.608 5.608	2.471 5.608 7.92	2 3 4	2 3 4	[150]
Deuterium <sup>b</sup>	D <sub>2</sub>	0.167	1.7 2.4	0.6 0.3	1.29 1.17	0.12 0.03	3.678 6.263	6.263 9.014	2 3	8 4	[63,144]
Helium <sup>a</sup>	He	0.123	0.674	0.011	1.366	0.002	2.47	9.39	2	3	[145]
Hydrogen <sup>b</sup>	H <sub>2</sub>	0.071	1.128 1.49 2.38	0.006 0.08 0.19	1.829 1.51 1.23	0.013 0.03 0.03	0 1.105 3.080	1.105 3.080 9.962	1 2 4	5 3 10	[63,144,215]
Hydrogen <sup>a</sup>	H <sub>2</sub>	0.089	1.80	0.12	1.89	0.09	0.801	1.525	2	4	[109]
Methane <sup>d</sup>	CH <sub>4</sub>	0.423	2.19 2.87	-- 0.09	1.35 1.166	-- 0.014	2.222 3.568	3.568 8.341	2 4	2 4	[150]
Nitrogen <sup>k</sup>	N <sub>2</sub>	0.0013	0.38	0.02	1.038	0.004	3.80	8.99	2	10	[57]
Nitrogen <sup>c</sup>	N <sub>2</sub>	0.811	0.94 1.14 2.1 4.0	0.10 0.18 0.3 0.2	1.83 1.59 1.26 0.88	0.09 0.08 0.06 0.04	0 1.51 3.26 5.2	1.57 3.26 5.23 8.63	1 2 3 4	9 15 24 12	[60,61,62,121,146, 149,179,213,233] / [149,160,221]
Oxygen <sup>c</sup>	O <sub>2</sub>	1.202	1.60 2.35	0.16 0.10	1.45 1.22	0.06 0.02	2.06 2.91	2.98 6.766	2 4	9 16	[121,146,223]
Xenon <sup>k</sup>	Xe	0.012	0.2 1.5 0.04 1.8	0.4 0.5 0.3 0.5	1.15 0.74 1.14 0.93	0.13 0.13 0.05 0.05	1.58 3.13 4.11 7.69	3.33 4.11 7.69 11.1	1 3 2 4	4 6 7 3	[71,81,216]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower Up (km/sec)	upper Up (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Xenon <sup>g</sup>	Xe	3.006	1.33	0.16	1.33	0.09	1.185	2.51	1	8	
			1.7	-	1.1	-	2.51	2.7	3	2	
			1.49	0.15	1.21	0.05	2.7	3.82	2	8	
			1.94	0.14	1.09	0.03	3.82	5.502	4	3	[151,212,213] /[161,212]
<b>Elements:</b>											
Antimony	Sb	6.695	3.2	-	-0.8	-	0	0.311	1	2	
			2.62	0.02	0.95	0.03	0.311	0.997	2	6	
			2.03	0.07	1.61	0.04	0.989	2.699	4	13	[121,126,217,227]
Bismuth	Bi	9.817	2.17	0.06	-1.0	0.5	0	0.32	1	17	
			1.08	0.06	2.20	0.07	0.32	1.183	2	30	
			2.01	0.04	1.358	0.019	1.183	4.45	4	21	[30,68,89,113,121, 131,166,217,226]
<b>Carbon:</b>											
Graphite	C	0.4665	0.4	0.3	1.14	0.06	2.114	6.147	2	6	[121]
Graphite	C	1.000	0.79	0.12	1.30	0.03	0.772	5.617	2	36	[121]
Graphite	C	1.611	1.75	0.09	1.42	0.05	0.911	4.22	2	60	[82,121,133,213, 217]
Graphite	C	1.794	2.04	0.14	1.66	0.08	0	2.563	2	30	
			4.2	0.5	0.71	0.18	2.372	3.08	3	19	
			1.9	0.3	1.49	0.07	3.069	5.42	4	41	[82,121,133,213, 217]
Graphite	C	2.205	3.11	0.07	4.7	0.2	0.012	0.41	1	12	
			4.19	0.05	1.83	0.04	0.404	1.9	2	77	
			7.5	0.3	0.21	0.11	1.89	3.316	3	24	
			3.92	0.06	1.331	0.008	3.119	28.38	4	22	[58,65,82,121,126, 133,162,217]
Diamond	C	1.90	1.2	0.2	1.73	0.05	2	6.5	2	5	[153]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Diamond	C	3.191	7.74	0.05	1.456	0.019	1.364	3.133	2	3	[121]
Diamond <sup>a</sup>	C	3.51	12.16	--	1	--	2	8.5	2	--	[153]
Glassy Carbon	C	1.507	2.72	0.11	1.12	0.03	0	5.8	2	45	[82,121,181]
Carbon Foam	C	0.435	0.85 -0.35	0.09 0.12	0.88 1.34	0.05 0.02	0.815 2.119	2.301 6.734	2 4	20 83	[121]
Carbon Fibers	C	1.519	1.18 2.52	0.09 0.10	1.73 1.14	0.06 0.03	0.924 2.361	2.361 5.041	2 4	5 15	[121]
Cobalt	Co	2.594	-0.15	0.06	1.602	0.017	0.651	6.39	2	11	[209]
Cobalt	Co	4.15	0.05	0.03	1.863	0.014	0.615	3.63	2	8	[209]
Cobalt	Co	5.533	0.42 1.38	0.04 0.08	2.11 1.76	0.02 0.02	0.293 2.89	2.89 5.2	2 4	10 4	[209]
Cobalt	Co	8.82	4.53 4.77 3.98	-- 0.02 0.13	1.77 1.285 1.66	-- 0.014 0.04	0 0.482 2.289	0.482 2.297 4.32	1 2 4	2 17 4	[24,121,126,131, 166,217,226]
Copper	Cu	1.909	0.03	0.08	1.361	0.009	0.661	26.1	2	27	[24,209]
Copper	Cu	2.887	0.37	0.08	1.406	0.015	1.15	17.25	2	28	[24,121,209]
Copper	Cu	3.57	0.03	0.02	1.675	0.008	0.63	3.96	2	6	[209]
Copper	Cu	4.475	1.35 0.15 1.9	-- 0.05 0.3	-2.02 1.87 1.33	-- 0.03 0.05	0 0.315 2.944	0.315 2.944 9.56	1 2 4	2 14 8	[56,121,134,136, 204,209]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower Up (km/sec)	upper Up (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Copper	Cu	6.144	2.73	0.11	-1.7	0.8	0	0.534	1	14	[56,121,128,134, 136,172,204,209, 217]
			0.87	0.07	1.97	0.04	0.534	3.365	2	39	
			3.2	0.4	1.27	0.07	3.327	8.77	4	5	
Copper	Cu	7.315	3.15	0.14	-0.4	0.2	0	0.701	1	3	[56,121,126,128, 134,136]
Copper	Cu	7.90	3.39	0.04	-0.06	0.08	0	0.646	1	3	[56,121,127,134, 136]
Copper	Cu	8.931	3.982	0.014	1.460	0.006	0	12.1	2	315	[22,24,25,28,29,30, 90,121,126,131, 134,136,139,141, 143,166,172,184, 205,213,217,226]
Germanium	Ge	5.328	5.93	0.19	-1.8	0.2	0.0775	1.226	1	17	[79,121,126,136, 213]
Gold	Au	19.263	2.95	0.03	1.81	0.07	0	0.71	1	5	[24,100,121,131, 166,217,226]
Indium	In	7.281	2.54	0.05	1.49	0.03	0.56	2.932	2	15	[24,121,166,217, 226]
Iodine	I <sub>2</sub>	4.902	1.62	0.03	1.25	0.04	0.49	0.9	1	5	[125,213]
	1.34	0.04	1.59	0.02	0.9	2.66	2	38			
	2.4	0.3	1.17	0.08	2.65	4.73	4	13			
Iridium	Ir	22.54	3.81	0.05	1.76	0.10	0	0.933	1	6	[24,121,136]
	4.37	0.07	1.15	0.05	0.933	1.629	3	4			
	3.36	0.09	1.76	0.04	1.629	3.09	4	3			

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Iron	Fe	2.633	-0.04 1.8	0.08 0.3	1.63 1.28	0.04 0.03	0.646 3.27	3.27 21.54	2 4	6 8	[204,209]
Iron	Fe	3.359	0.23	0.09	1.67	0.04	0.644	5.52	2	24	[121,134,136,204]
Iron	Fe	4.547	0.57 2.4	0.09 0.3	1.88 1.38	0.04 0.05	0.591 3.58	3.59 9.1	2 4	33 6	[121,134,136,204, 209]
Iron	Fe	5.783	3.15 1.17 0.01 2.25	0.10 0.03 0.09 0.09	-1.7 1.98 3.22 1.61	0.6 0.04 0.08 0.03	0 0.537 0.941 1.565	0.537 0.941 1.249 4.95	1 2 3 4	12 4 3 10	[31,121,134,136, 172,213]
Iron	Fe	6.972	4.1 2.4 1.2 2.77	-- 0.2 0.3 0.08	-1.5 1.3 2.8 1.71	-- 0.3 0.2 0.04	0 0.569 0.85 1.453 3.131	0.569 0.85 1.453 3.131	1 2 3 4	2 5 4 15	[121,134,136]
Iron	Fe	7.853	5.85 3.48 3.94 5.36	0.12 0.05 0.03 0.07	-1.7 1.91 1.584 1.302	0.8 0.05 0.013 0.008	0 0.763 1.413 4.50	0.573 1.433 4.55 21.73	1 2 4a 4b	16 42 97 18	[24,25,29,30,31,33, 108,121,126,131, 132,134,136,138, 162,166,172,202, 206,213,217,226] / [6,20,43,44]
Iron-Nickel (see Taenite)	(Fe, Ni)										
Iron-Silicon	Fe <sub>12</sub> Si	7.641	3.87	0.04	1.67	0.02	0.984	3.568	2	37	[42,121]
Iron-Silicon	Fe <sub>7</sub> Si	7.49	4.01	0.06	1.71	0.04	0.975	2.291	2	3	[121]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower Up (km/sec)	upper Up (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Iron-Silicon (see Suessite)											
Lead	Pb	4.71	0.31 0.17 0.78	-- 0.06 0.12	1.42 1.59 1.37	-- 0.03 0.03	0.607 1.016 2.83	1.016 2.83 5.77	2 3 4	2 5 3	[209]
Lead	Pb	6.79	0.555 0.94 1.36	0.016 0.05 --	1.726 1.462 1.33	0.015 0.018 --	0.55 1.44 3.24	1.44 3.24 4.85	2 3 4	4 3 2	[209]
Lead	Pb	8.40	0.71 1.15 1.84	0.07 0.08 0.05	2.10 1.59 1.306	0.14 0.05 0.016	0.26 0.73 2.18	0.73 2.18 4.91	2 3 4	3 6 5	[209]
Lead	Pb	9.51	1.20 1.58	0.07 0.05	1.87 1.51	0.08 0.02	0.46 1.16	1.16 2.73	2 3	3 3	[209]
Lead	Pb	11.345	1.992 2.70	0.014 0.04	1.511 1.213	0.012 0.006	0 2.335	2.36 19.12	2 4	93 42	[25,28,30,33,121, 131,141,163,166, 206,213,217,226]
Mercury	Hg	13.54	1.45 1.752	-- 0.007	2.26 1.724	-- 0.009	0 0.56	0.56 0.991	2 4	2 3	[121,225]
Nickel	Ni	1.644	-0.04 -0.4	0.05 0.3	1.32 1.47	0.02 0.06	0.67 2.86	3.15 5.73	1 2	16 19	[209]
Nickel	Ni	3.202	-0.09 0.87	0.07 0.17	1.71 1.46	0.02 0.02	1.53 4.045	4.045 10.31	2 4	6 3	[209]
Nickel	Ni	4.198	0.02	0.13	1.88	0.05	0.61	3.86	2	15	[209]
Nickel	Ni	5.15	0.7 1.40	-- 0.09	1.9 1.606	-- 0.019	1.23 1.98	1.98 8.91	2 4	2 7	[209]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Nickel	Ni	6.275	0.23	0.09	2.79	0.13	0.54	0.80	1	3	
			1.02	0.10	2.03	0.05	0.80	3.28	2	9	
			2.15	--	1.64	--	3.28	4.62	4	2	[209]
Nickel	Ni	8.896	4.57	0.04	0.29	0.17	0	0.354	1	4	
			3.83	0.17	2.5	0.3	0.349	0.635	2	28	
			4.31	0.04	1.63	0.03	0.635	2.63	4a	52	
			5.41	0.08	1.300	0.015	2.63	7.5	4b	18	[24,25,90,121,131,166,209,213,217,226]
Palladium	Pd	11.996	3.83	0.02	1.83	0.04	0.00	0.825	2	7	
			4.09	0.05	1.49	0.03	0.803	2.317	4	13	[56,121,136,166,217,226]
Platinum	Pt	21.445	3.587	0.014	1.556	0.008	0.00	3.444	2	29	[86,121,136,166,217,226]
Rhenium	Re	20.53	4.12	0.05	-0.04	0.18	0.00	0.372	1	3	
			3.56	0.08	1.63	0.08	0.372	1.441	2	7	
			4.0	0.2	1.32	0.12	1.346	2.028	4	6	[121,136]
Rhenium	Re	20.984	4.16	0.04	1.40	0.06	0.00	1.127	2	7	[121,136]
Rhodium	Rh	12.422	4.28	0.12	2.7	0.4	0.00	0.426	1	4	
			4.76	0.05	1.41	0.04	0.369	2.004	2	14	
			4.043	0.018	1.713	0.006	2.004	3.8	4	3	[24,121,136,166,226]
Silver	Ag	10.49	3.23	0.04	1.59	0.03	0.00	2.149	2	16	
			3.56	0.13	1.46	0.04	2.12	4.32	4	9	[24,30,56,121,131,166,217,226]
Suessite	(Fe,Ni) <sub>3</sub> Si	6.870	5.21	--	2.25	--	0.00	0.495	1	2	
			5.53	0.06	1.23	0.03	0.495	3.627	2	32	[42,121]
Sulfur	S	2.02	3.633	0.013	0.606	0.010	0.897	1.470	2	3	
			2.8	0.3	1.18	0.15	1.431	2.046	4	6	[121]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower Up (km/sec)	upper Up (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Taenite (also Kamacite)	(Fe,Ni)	7.933	4.41	0.05	1.01	0.05	0.00	1.09	1	11	[36,121,132,136, 217]
			3.79	0.051	1.65	0.02	1.019	2.777	2	41	
			4.20	0.17	1.48	0.05	2.723	4.59	4	11	
Tin	Sn	7.299	2.60	0.15	2.2	0.9	0.00	0.304	1	3	[24,25,30,121,126, 131,166,213,217, 226]
			3.33	0.07	-0.14	0.15	0.304	0.5	3	6	
			2.48	0.03	1.57	0.03	0.5	2.15	2	66	
			3.43	0.03	1.205	0.008	2.15	8	4	33	
Wairauite	CoFe	8.091	4.64	0.04	1.63	0.08	0.00	0.647	2	12	[121]
		5.69	0.02	-0.10	0.02	0.02	0.663	1.037	3	5	
		3.78	0.08	1.62	0.04	0.04	1.038	2.801	4	10	
Zinc	Zn	6.51	3.69	0.17	0.98	0.12	0.54	2.08	2	3	[23]
		3.04	0.15	1.35	0.05	0.05	2.08	5.04	4	4	
Zinc	Zn	7.138	3.00	0.02	1.586	0.013	0.00	3.01	2	39	[24,25,30,121,126, 131,166,217,224, 226]
		3.70	0.15	1.37	0.04	0.04	2.98	4.85	3	10	
		4.05	0.02	1.303	0.003	0.003	4.85	8	4	9	
<b>Carbides:</b>											
Moissanite	SiC	2.333	2.3	0.3	1.84	0.12	2.048	3.444	4	10	[121,127,136]
Moissanite	SiC	3.029	8.4	0.6	0.3	0.3	1.535	2.112	3	3	[121]
		5.6	0.3	1.62	0.14	0.14	2.112	2.842	4	4	
Moissanite	SiC	3.122	8.0	--	6.0	--	0.00	0.464	1	2	[121,127,136]
		10.29	0.13	-0.38	0.10	0.10	0.674	1.678	3	9	
		7.84	0.11	1.03	0.05	0.05	1.678	2.912	4	10	

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower Up (km/sec)	upper Up (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Tantalum Carbide	TaC	12.626	3.32	0.09	1.49	0.05	0.887	2.619	2	20	[121]
Tantalum Carbide	TaC	14.110	4.34	0.05	1.36	0.03	0.435	3.76	2	21	[121,152]
Tungsten Carbide <sup>P</sup>	WC	15.013	4.97 5.21	0.11 0.03	2.1 1.14	0.4 0.02	0.00 0.369	0.369 1.819	1 2	4 12	[121,127,136]
Tungsten Carbide	WC	15.66	4.926	0.014	1.163	0.007	0.45	3.66	2	4	[152]
<b>Sulfides:</b>											
Sphalerite	ZnS	3.952	5.08 3.10 0.9	0.03 0.10 --	-0.09 1.22 2.0	0.03 0.04 --	0.63 1.52 2.70	1.52 2.70 3.56	2 3 4	3 3 2	[186]
Pyrhotite	Fe <sub>1-x</sub> S	4.605	5.8 2.31 3.23	0.2 0.17 0.10	-4.7 2.08 1.49	0.5 0.15 0.03	0.235 0.494 1.496	0.547 1.599 5.361	1 2 4	3 10 14	[3,53]
Pyrite	FeS <sub>2</sub>	4.933	8.8 5.3	1.0 0.10	-1.4 1.47	1.0 0.04	0.225 1.133	1.39 5	1 2	7 11	[10,186]
Potassium Iron Sulfide	KFeS <sub>2</sub>	2.59	2.32 8.2 0.25	0.06 -- 0.07	1.97 -1.0 1.912	0.05 -- 0.017	0.223 2.05 2.79	2.05 2.79 4.72	2 3 4	11 2 3	[191,230]
<b>Halides:</b>											
Griceite	LiF	1.27	0.74	--	1.58	--	2.4	6.59	2	2	[106]
Griceite	LiF	2.638	5.10	0.09	1.35	0.03	0.452	10.01	2	68	[38,54,121,195,213] /[177]
Villiaumite	NaF	2.792	4.08 6.7 2.83	0.14 0.3 0.04	1.42 -0.28 1.635	0.13 0.16 0.013	0.5 1.54 2.027	1.54 2.027 3.982	2 3 4	8 4 8	[54,213]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Halite	NaCl	0.868	-0.15	0.19	1.72	0.06	1.942	3.62	2a	4	[121]
			7.0	1.3	-0.2	0.3	3.62	4.181	3	3	
			-0.4	--	1.5	--	4.181	5.552	2b	2	
Halite	NaCl	0.989	0.74	0.09	1.470	0.018	2.53	6.02	2	5	[106]
			3.9	--	0.9	--	6.02	6.6	3	2	
			-17.0	--	4.1	--	6.6	6.7	4	2	
Halite	NaCl	1.427	1.56	0.12	1.49	0.03	2.29	5.66	2	5	[106]
			5.6	--	0.8	--	5.66	6	3	2	
			-5.1	--	2.5	--	6	6.11	4	2	
Halite	NaCl	2.159	3.60	0.09	1.17	0.18	0.00	0.647	1	28	[32,73,84,106,121, 213]/[48, 103, 107, 176, 177]
			3.41	0.03	1.42	0.03	0.646	1.7	2a	113	
			4.35	0.09	0.88	0.04	1.7	2.5	3a	109	
			2.43	0.07	1.66	0.02	2.5	3.75	4a	95	
			4.4	0.3	1.11	0.08	3.73	4.356	3b	43	
			3.5	0.3	1.33	0.05	4.324	6.52	4b	8	
			19	--	-1	--	6.52	6.8	3c	2	
			3.8	0.2	1.18	0.02	6.8	11.05	4c	3	
Sylvite	KCl	0.79	0.9	--	1.3	--	2.66	7.19	2	2	[106]
Sylvite	KCl	1.41	1.9	--	1.3	--	2.3	6.56	2	2	[106]
Sylvite	KCl	1.986	2.86	0.09	1.26	0.09	0.249	2.2	1	33	[34,38,85,106,213] /[48,107,177]
			4.0	0.2	1.09	0.05	2.2	6.71	2	9	
			14	--	-0.4	--	6.71	7.1	3	2	
			2.5	0.4	1.25	0.04	7.1	11.38	4	3	
Potassium Bromide	KBr	2.747	2.83	0.16	-0.1	0.3	0.27	0.61	1	4	[38,106,121,213] / [48]
			1.88	0.05	1.50	0.03	0.57	2.9	2	13	
			2.63	0.09	1.24	0.02	2.862	5.09	4a	12	
			3.23	0.19	1.11	0.03	5.09	10.6	4b	6	

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower Up (km/sec)	upper Up (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Cesium Iodide	CsI	2.51	0.66 1.37	0.12 0.12	1.42 1.22	0.04 0.02	1.04 4.23	4.23 6.75	2 4	6 4	[154]
Cesium Iodide	CsI	4.51	1.57	0.17	1.66	0.17	0.56	1.32	2	7	
			3.8	--	0.1	--	1.32	1.56	3	2	
			1.95	0.05	1.302	0.019	1.56	4.3	4a	13	
			2.66	0.13	1.141	0.019	4.3	9.28	4b	3	[38,121,154,213] /[199,209]
Fluorite	CaF <sub>2</sub>	3.18	5.5	--	0.8	--	1.1	2.22	1	2	
			4.64	0.16	1.19	0.06	2.22	3.38	2	4	
			8.2	--	0.14	--	3.38	3.67	3	2	
			0.4	0.3	2.27	0.07	3.67	5.76	4	5	[35]
Cryolite	Na <sub>3</sub> AlF <sub>6</sub>	2.96	4.70	0.10	0.89	0.09	0.71	1.57	2	3	
			3.76	0.12	1.44	0.04	1.57	3.8	4	4	[186]
<b>Oxides:</b>											
Water, Ice <sup>j</sup>	H <sub>2</sub> O	0.35	0.080	0.18	1.40	0.03	2.76	6.75	2	5	[41]
Water, Ice <sup>j</sup>	H <sub>2</sub> O	0.60	0.83	0.16	1.40	0.03	2.57	6.2	2	5	[41]
Water, Ice <sup>j</sup>	H <sub>2</sub> O	0.915	4.05	0.05	-1.89	0.16	0.045	0.858	1	7	
			1.43	0.11	1.48	0.03	0.858	5.67	2	9	[41,75,114]
Water, Ice <sup>k</sup>	H <sub>2</sub> O	0.999	1.47	0.04	1.93	0.06	0.00	0.97	1	14	
			1.70	0.06	1.71	0.03	0.9	2.53	2	58	
			2.64	0.07	1.270	0.008	2.479	32.42	4	25	[19,29,116,121,140, 157,167,213,217, 225] / [87,119]
Seawater	(mixture)	1.03	1.69 2.07	0.08 0.09	1.73 1.38	0.10 0.03	0.31 1.11	1.11 4.76	2 4	3 4	[210]
Bromellite	BeO	2.454	3.5	0.3	1.92	0.12	1.799	3.356	2	6	[120,121]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Bromellite	BeO	2.661	8.5 5.4	-- 0.8 †	-1.2 1.6	-- 0.4	0.577 1.25	1.25 3.46	1 2	2 3	[120,121,213]
Bromellite	BeO	2.797	9.10 6.7	0.18 0.5	-1.0 1.32	0.2 0.18	0.368 1.094	1.094 3.71	1 2	4 5	[120,121,213]
Bromellite	BeO	2.886	7.76 8.42	0.10 0.06	1.22 1.042	0.05 0.013	0.77 2.74	2.74 5.78	2 4	12 3	[120,121,152,213]
Bromellite	BeO	2.989	10.84 9.65 7.7	0.08 0.10 0.3	-0.78 0.49 1.53	0.13 0.07 0.11	0.317 0.939 1.825	0.939 1.825 2.822	1 2 4	4 4 4	[120,121]
Periclase	MgO	2.842	2.8	0.2	1.84	0.08	1.749	3.528	2	6	[56,121,127]
Periclase	MgO	3.00	3.10	0.14	1.88	0.06	1.259	3.362	2	13	[56,121,127]
Periclase	MgO	3.355	3.5 4.78 5.73	0.3 0.13 0.12	2.7 1.77 1.36	0.3 0.06 0.03	0.629 1.509 2.619	1.557 2.652 5.62	2a 2b 4	8 9 11	[37,56,121,127]
Periclase	MgO	3.583	6.09 6.83	0.10 0.12	1.75 1.29	0.08 0.04	0.626 1.92	1.967 4.44	2 4	25 16	[2,47,56,76,121, 127,217,218] / [197]
Magnesio-wüstite	Mg <sub>0.6</sub> Fe <sub>0.4</sub> O	4.397	4.81 6.50 4.6	0.04 0.16 0.7	1.63 0.91 1.51	0.02 0.05 0.17	1.55 2.3 3.33	2.3 3.33 4.29	2 3b 4	3 4 4	[219]
Magnesio-wüstite	Mg <sub>0.1</sub> Fe <sub>0.9</sub> O	5.191	4.3 6.27 4.96 5.53 4.76	-- 0.11 0.07 0.12 0.14	5.7 -0.3 1.56 1.14 1.49	-- 0.2 0.07 0.07 0.06	0.00 0.332 0.705 1.28 2.169	0.332 0.705 1.28 2.169 2.721	1 3a 2 3b 4	2 3 3 4 3	[121]
Lime	CaO	2.980	3.56 6.7 2.61	0.11 0.6 0.16	1.79 0.3 1.67	0.07 0.2 0.05	0.882 2.083 3.01	2.083 3.01 3.558	2 3 4	6 3 5	[121]



**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Lime	CaO	3.324	7.4	0.8	0.5	0.3	1.801	3.285	3	5	[97] / [50]
Wüstite	Fe <sub>1-x</sub> O	5.548	4.80	0.10	1.33	0.07	1.22	1.766	2	3	
		6.9	--	0.2	--	--	1.766	2.034	3	2	
		3.72	0.15	1.59	0.04	--	2.414	4.055	4	4	[97]
Wüstite	Fe <sub>1-x</sub> O	5.662	3.4	--	2.4	--	1.56	1.73	2	2	
		6.84	0.02	0.408	0.011	--	1.73	2.51	3	3	
		3.3	--	1.8	--	--	2.51	2.62	4	2	[232]
Corundum	Al <sub>2</sub> O <sub>3</sub>	3.761	11.04	0.18	-4.3	0.5	0.18	0.706	1	7	
		6.61	0.16	1.35	0.07	--	0.706	3.282	2	15	[9,121,127]
Corundum	Al <sub>2</sub> O <sub>3</sub>	3.843	10.20	0.125	-1.9	0.3	0.18	0.898	1	5	
		7.08	0.11	1.36	0.05	--	0.898	2.979	2	10	[47,121,122,127, 217]
Corundum	Al <sub>2</sub> O <sub>3</sub>	3.92	8.71	0.05	0.716	0.017	0.37	5.5	2	6	[152]
Corundum	Al <sub>2</sub> O <sub>3</sub>	3.979	11.04	0.07	1.1	0.3	0.033	0.46	0	41	
		17.8	1.6	-14	3	0.46	0.621	1	6		
		8.83	0.06	0.93	0.03	0.555	3.064	2	52	[9,47,78,121,122, 127,217]	
Corundum	Al <sub>2</sub> O <sub>3</sub>	4.00	9.52	0.04	0.955	0.008	1.02	8.28	2	4	[152]
Hematite	α-Fe <sub>2</sub> O <sub>3</sub>	5.047	6.18	0.12	1.40	0.17	0.00	1.03	2	6	
		7.435	0.011	0.035	0.007	--	1.097	2.294	3	4	
		4.39	0.11	1.37	0.04	--	2.294	3.18	4	10	[47,121,127,200, 217]
Ilmenite	Fe <sup>+2</sup> TiO <sub>3</sub>	4.75	5.85	0.08	1.28	0.07	0.85	1.38	2	3	
		7.43	--	0.13	--	--	1.38	1.92	3	2	
		5.46	--	1.15	--	--	1.92	3.09	4	2	[186]
Ilmenite	Fe <sup>+2</sup> TiO <sub>3</sub>	4.787	6.33	0.12	1.0	0.2	0.00	0.652	2	3	

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower Up (km/sec)	upper Up (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Perovskite	CaTiO <sub>3</sub>	3.86	6.86	0.06	0.17	0.05	0.626	2.009	3	5	[121,127]
			4.07	0.18	1.54	0.07	2.009	3.082	4	12	
			5.25	0.11	1.48	0.07	0.59	2.35	2	5	
Barium Titanate	BaTiO <sub>3</sub>	5.447	6.5	0.3	0.93	0.10	2.35	3.24	3	3	[187]
			4.0	--	1.7	--	3.24	5.42	4	2	
			2.33	0.10	2.63	0.10	0.533	1.334	2	23	
Spinel	MgAl <sub>2</sub> O <sub>4</sub>	2.991	3.7	0.2	1.60	0.12	1.321	2.479	4	9	[66,121,213]
			4.05	0.19	1.43	0.07	1.727	3.63	4	23	
			7.1	--	1.13	--	0.00	0.987	2	2	
Spinel	MgAl <sub>2</sub> O <sub>4</sub>	3.417	8.23	0.06	-0.03	0.03	0.987	2.146	3	10	[47,121,127,217]
			5.07	0.16	1.42	0.06	2.134	3.348	4	13	
			8.04	0.07	0.27	0.04	0.688	2.311	3	5	
Spinel	MgAl <sub>2</sub> O <sub>4</sub>	3.514	7.26	0.07	1.48	0.17	0.00	0.688	2	3	[121,127]
			5.7	0.5	1.35	0.16	2.311	3.507	4	8	
			7.2	--	-0.2	--	1.14	1.5	3	2	
Magnetite	Fe <sup>+2</sup> Fe <sup>+3</sup> <sub>2</sub> O <sub>4</sub>	5.07	3.0	1.4	1.8	0.4	2.72	4.52	4	3	[186]
			5.9	--	1.3	--	0.00	0.61	2	2	
			6.56	0.08	0.05	0.06	0.61	1.786	3	10	
Rutile	TiO <sub>2</sub>	4.21	4.24	0.11	1.36	0.05	1.757	2.975	4	15	[47,121,127,217]
			6.96	--	0.23	0.03	1.14	2.44	3	3	
			2.1	0.3	2.15	0.08	2.44	5.2	4	5	
Rutile	TiO <sub>2</sub>	4.245	10.3	0.5	-4.3	1.3	0.09	0.676	1	11	[11,47,121,123,127, 130,217]
			7.68	0.11	0.21	0.07	0.468	2.858	3	8	
			3.0	0.6	1.8	0.2	2.858	3.191	4	8	
Pyrolusite	Mn <sup>+4</sup> O <sub>2</sub>	4.318	3.77	0.14	1.46	0.07	0.769	3.263	2	16	[47,121,217]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Cassiterite	SnO <sub>2</sub>	6.694	6.82	0.05	-0.22	0.05	0.509	1.866	1	7	
			5.15	0.06	0.68	0.03	1.866	2.501	2	4	
			2.6	0.6	1.7	0.2	2.501	2.833	4	3	[47,121,217]
Argutite	GeO <sub>2</sub>	6.277	9.51	0.12	0.29	0.11	0.15	2.57	2	6	[92]
Baddeleyite	ZrO <sub>2</sub>	4.512	4.4	--	0.27	--	0.00	1.622	2	2	
Baddeleyite	ZrO <sub>2</sub>	4.512	1.9	0.4	1.88	0.17	1.622	2.994	4	8	[121]
			5.814	5.17	0.08	1.02	0.05	0.41	2.17	2	4
			4.42	0.07	1.35	0.03	2.17	2.99	4	4	[124]
Cerianite	(Ce <sup>+4</sup> ,Th)O <sub>2</sub>	1.133	0.2	0.3	1.20	0.07	1.925	5.437	2	7	[121]
Uraninite	UO <sub>2</sub>	10.337	3.99	0.06	0.20	0.13	0.00	0.571	1	3	
			3.59	0.04	0.91	0.03	0.568	1.983	2	12	
			1.7	0.4	1.8	0.16	1.983	2.493	4	6	[121]
Uraninite	UO <sub>2</sub>	6.347	0.43	0.07	1.70	0.03	1.025	3.286	4	18	[121]
Uraninite	UO <sub>2</sub>	4.306	0.12	0.11	1.51	0.04	0.88	3.855	4	15	[121]
Uraninite	UO <sub>2</sub>	3.111	-0.22	0.10	1.47	0.03	1.355	4.256	4	15	[121]
<b>Hydroxides:</b>											
Brucite	Mg(OH) <sub>2</sub>	2.37	4.75	0.06	1.26	0.02	1.25	3.41	2	6	
Brucite	Mg(OH) <sub>2</sub>	2.383	0.9	--	2.4	--	3.41	3.96	4	2	[186]
			5.0	0.2	1.22	0.11	0.886	3.079	2	13	[69]
Goethite	$\alpha\text{-Fe}^{+3}\text{O(OH)}$	4.0	4.4	--	1.6	--	1.02	1.34	2	2	
			5.77	0.11	0.61	0.06	1.34	2.52	3	3	
			2.9	--	1.8	--	2.52	3.51	4	2	[187]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
<b>Carbonates:</b>											
Calcite	CaCO <sub>3</sub>	2.701	6.9 3.71	0.2 0.03	-2.8 1.435	1.0 0.013	0.081 0.81	0.81 3.845	1 2	24 6	[11,21,102] / [103]
Magnesite	MgCO <sub>3</sub>	2.975	6.08	0.09	1.26	0.04	0.6	3.61	2	6	[102]
Dolomite	CaMg(CO <sub>3</sub> ) <sub>2</sub>	2.828	6.2 5.30	0.5 0.10	0.4 1.16	0.5 0.03	0.495 1.12	1.15 5.32	2 4	5 19	[102,184,213]
Aragonite	CaCO <sub>3</sub>	2.928	5.82	0.11	0.78	0.12	0.11	1.83	2	12	[220]
<b>Sulfates:</b>											
Mascagnite	(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	1.3	0.77 1.8	0.15 0.2	2.28 1.54	0.167 0.12	0.36 1.15	1.21 2.5	2 4	8 11	[99]
Mascagnite	(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	1.6	1.96	0.10	2.09	0.10	0.28	1.83	2	8	[99]
Mascagnite	(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	1.73	3.71	0.14	1.34	0.12	0.2	1.87	2	5	[99]
Anhydrite	CaSO <sub>4</sub>	2.97	3.60 4.6 3.24	0.06 -- 0.11	1.75 1.1 1.72	0.05 -- 0.03	0.73 1.55 2.42	1.55 1.85 3.71	2 3 4	3 2 4	[186]
Barite	BaSO <sub>4</sub>	4.375	3.3 4.7 2.3	-- 0.2 0.4	1.9 0.54 1.86	-- 0.15 0.16	0.64 1.03 1.69	1.03 1.69 3.29	2 3 4	2 3 5	[186]
Gypsum	CaSO <sub>4</sub> · 2H <sub>2</sub> O	2.28	2.80 5.2 2.49	0.17 -- 0.12	1.95 0.5 1.79	0.13 -- 0.04	0.85 1.72 2.15	1.72 2.15 4.06	2 3 4	3 2 5	[186] / [103]
<b>Borates:</b>											
Sassolite	H <sub>3</sub> BO <sub>3</sub>	1.471	2.09 1.14	0.11 --	1.27 1.85	0.08 --	1.254 1.639	1.639 2.114	2 4	3 2	[121]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
<b>Silica Polymorphs:</b>											
Quartz	SiO <sub>2</sub>	2.651	6.61 5.65 8.14 6.44 5.29 1.48 8.2 4.0	0.12 0.09 0.11 -0.43 0.08 0.10 0.33 0.2	1.02 0.9 -1.32 0.08 0.20 1.80 0.20 1.283	0.19 0.2 0.12 0.08 0.04 0.03 0.12 0.018	0.295 0.285 0.508 0.48 1.803 2.46 4.51 4.84	0.81 0.66 1.815 1.815 2.48 4.55 4.84 26.76	0a 0b 1a 1b 3a 2 3b 4	16 29 16 12 6 24 3 10	[11,17,27,37,47,72, 118,121,127,129, 155,162,208,217, 222] / [118,165]
Porous Quartz	SiO <sub>2</sub>	1.15	0.41	0.09	1.40	0.03	1.52	5.28	2	15	[188,207]
Porous Quartz	SiO <sub>2</sub>	1.43	0.87	0.09	1.38	0.03	0.65	5.08	2	30	[188,207]
Porous Quartz	SiO <sub>2</sub>	1.766	1.27 3.42 0.25	0.03 0.13 0.05	1.356 0.34 1.589	0.019 0.05 0.011	0.62 2.13 2.55	2.13 2.55 9.0	2 3 4	4 3 9	[188,207]
Porous Quartz	SiO <sub>2</sub>	1.877	2.75 -0.5	0.07 0.4	0.82 1.86	0.03 0.10	0.849 3.374	3.374 4.863	2 4	11 15	[121]
Porous Quartz	SiO <sub>2</sub>	2.151	3.02 0.7	0.16 0.2	0.84 1.67	0.08 0.06	0.799 3.199	3.199 6.52	2 4	11 16	[121,207]
Silicic Acid	H <sub>4</sub> SiO <sub>4</sub>	0.55	-0.09	0.11	1.27	0.02	3.51	6.59	4	5	[188]
Silicic Acid	H <sub>4</sub> SiO <sub>4</sub>	0.65	-0.01	0.03	1.275	0.006	2.94	5.93	4	6	[188]
Silicic Acid	H <sub>4</sub> SiO <sub>4</sub>	0.80	0.78 0.33	0.03 0.04	1.05 1.261	0.02 0.009	0.68 1.98	1.98 5.71	2 4	3 7	[188]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Aerogel	SiO <sub>2</sub> <sup>r</sup>	0.172	0.21	0.07	1.01	0.02	1.494	4.01	2	6	
			-0.44	0.10	1.164	0.018	4.01	7.401	4	8	[88,188]
Aerogel	SiO <sub>2</sub> <sup>r</sup>	0.295	0.21	0.03	1.11	0.05	0.302	0.765	2	3	[158]
Aerogel	SiO <sub>2</sub> <sup>r</sup>	0.40	0.29	0.03	1.089	0.010	1.456	3.63	2	4	
			-0.214	0.013	1.224	0.003	3.63	6.37	4	4	[188]
Aerogel	SiO <sub>2</sub> <sup>r</sup>	0.55	0.590	0.006	0.945	0.006	0.335	1.43	2	3	
			0.15	0.04	1.219	0.008	1.43	6.1	4	7	[188]
Cristobalite	SiO <sub>2</sub>	2.13	2.29	0.09	1.01	0.06	0.96	1.99	2	3	
			1.23	0.05	1.545	0.014	1.99	4.66	4	6	[156]
Coesite	SiO <sub>2</sub>	1.15	0.44	0.07	1.44	0.02	1.2	4.59	2	5	[156]
Coesite	SiO <sub>2</sub>	2.40	3.4	0.3	1.42	0.19	1.33	2.05	2	3	
			4.3	--	0.97	--	2.05	4.16	3	2	[156]
Coesite	SiO <sub>2</sub>	2.92	5.833	0.010	0.902	0.009	0.66	1.48	2	3	
			6.92	0.02	0.168	0.011	1.48	2.86	3	3	
			2.6	0.4	1.68	0.12	2.86	4.05	4	3	[156]
<b>Silicates:</b>											
Forsterite	Mg <sub>2</sub> SiO <sub>4</sub>	3.059	5.71	0.07	0.64	0.03	0.825	2.771	2	6	
			3.0	0.3	1.65	0.10	2.771	3.626	4	6	[47,121,127,217]
Forsterite	Mg <sub>2</sub> SiO <sub>4</sub>	3.212	10.5	0.2	-3.8	0.4	0.18	0.797	1a	16	
			6.8	0.3	1.4	0.6	0.00	0.795	1b	12	
			7.21	0.10	0.55	0.06	0.749	2.449	2	20	
			4.6	0.3	1.51	0.08	2.433	4.61	4	20	[91,121,127,201, 229] / [165]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Olivine	(Mg,Fe) <sub>2</sub> -SiO <sub>4</sub> : (Mg <sub>0.92</sub> -Fe <sub>0.08</sub> ) <sub>2</sub> SiO <sub>4</sub>	3.264	8.84	0.13	-0.99	0.18	0.272	1.43	1	4	
			5.6	--	1.3	--	1.43	2.19	2	2	
			8.1	--	0.15	--	2.19	2.8	3	2	
			6.0	0.3	0.88	0.08	2.8	4.81	4	3	[15,121]
Fayalite	Fe <sub>2</sub> SiO <sub>4</sub>	4.245	6.17	0.09	0.23	0.06	0.702	1.967	2	10	
			3.78	0.13	1.42	0.05	1.967	3.483	4	10	[47,121,127,217]
Zircon	ZrSiO <sub>4</sub>	4.549	8.58	0.18	-1.3	0.4	0.11	1.07	1	7	
			7.14	0.05	0.02	0.03	1.07	2.52	3	4	
			-1.6	0.9	3.5	0.3	2.52	2.84	2	4	[124]
Almandine	(Fe <sub>0.79</sub> -Mg <sub>0.14</sub> -Ca <sub>0.04</sub> -Mn <sub>0.03</sub> ) <sub>3</sub> -Al <sub>2</sub> Si <sub>3</sub> O <sub>12</sub>	4.181	-10	--	50	--	0.29	0.32	1	2	
			5.92	0.07	1.39	0.08	0.45	1.14	2	8	
			3.0	--	3.6	--	1.29	1.46	3	2	
			6.4	0.3	1.33	0.16	1.46	1.8	4	7	[77]
Grossular	Ca <sub>3</sub> Al <sub>2</sub> Si <sub>3</sub> O <sub>12</sub>	3.45	8.3	0.17	0.47	0.10	0.18	3.04	2	16	[121]
Mullite	Al <sub>6</sub> Si <sub>2</sub> O <sub>13</sub>	2.668	2.30	0.13	1.65	0.04	1.935	4.077	2	13	[121,127]
Mullite	Al <sub>6</sub> Si <sub>2</sub> O <sub>13</sub>	3.154	8.732	0.016	-0.394	0.015	0.717	1.479	1	3	
			8.29	0.04	-0.09	0.02	1.479	2.003	3	3	
			6.5	--	0.78	--	2.003	3.311	4	2	[121]
Kyanite	Al <sub>2</sub> SiO <sub>5</sub>	2.921	7.45	0.06	-0.58	0.07	0.608	1.157	1	3	
			7.02	0.09	-0.19	0.05	1.157	2.359	3	5	
			2.2	0.3	1.85	0.10	2.359	3.383	4	5	[121]
Kyanite	Al <sub>2</sub> SiO <sub>5</sub>	3.645	7.8	--	0.6	--	1.537	2.745	2	2	
			3.9	1.7	2.0	0.6	2.745	3.22	4	3	[121]
Andalusite	Al <sub>2</sub> SiO <sub>5</sub>	3.074	5.3	--	1.9	--	0.00	1.1	2	2	
			6.92	0.15	0.31	0.07	1.1	2.817	3	12	
			2.9	0.4	1.80	0.13	2.817	3.73	4	8	[47,121,217]

TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Sillimanite	$\text{Al}_2\text{SiO}_5$	3.127	6.97	0.15	0.68	0.16	0.00	1.262	1	3	
			7.8	0.2	-0.10	0.11	1.068	2.461	3	8	
			3.8	0.2	1.57	0.08	2.461	3.611	4	10	[47,121,217]
Topaz	$\text{Al}_2\text{SiO}_4 \cdot (\text{F},\text{OH})_2$	3.53	8.10	0.09	0.054	0.08	0.052	1.75	1	4	
			5.3	0.4	1.722	0.15	1.75	3.59	2	6	[186]
Tourmaline	$\text{Ca}(\text{Al},\text{Fe},-\text{Mg})_3\text{Al}_6\text{Si}_6\text{O}_{18}(\text{OH},\text{F})_4$	3.179	6.2	--	1.2	--	0.824	1.555	1	2	
			8.1	0.4	0.05	0.15	1.555	2.888	3	7	
			3.6	0.5	1.62	0.16	2.888	3.695	4	8	[121]
Muscovite	$\text{KAl}_2(\text{Si}_3\text{Al})\text{O}_{10}(\text{OH},\text{F})_2$	2.835	3.3	0.3	1.95	0.16	1.27	2.44	2	3	
			6.2	--	0.7	--	2.44	3.18	3	2	
			4.5	0.2	1.29	0.06	3.18	4.74	4	4	[182]
Serpentine	$\text{Mg}_3\text{Si}_2\text{O}_5(\text{OH})_4$	2.621	5.30	0.15	0.90	0.11	0.431	2.025	2	10	
			6.5	0.4	0.20	0.18	1.719	2.561	3	10	
			3.8	0.5	1.34	0.12	2.658	5.427	4	16	[47,121,211,217]
<i>Pyroxenes:</i>											
Enstatite	$\text{Mg}_2\text{Si}_2\text{O}_6$	2.714	2.70	0.11	1.31	0.04	1.901	3.258	2	5	[47,121,217]
Enstatite	$\text{Mg}_2\text{Si}_2\text{O}_6$	2.814	2.74	0.14	2.04	0.09	0.746	1.956	2	5	
			6.8	--	-0.1	--	1.956	2.128	3	2	
			3.64	0.16	1.43	0.05	2.128	3.946	4	9	[121,127]
Enstatite	$\text{Mg}_2\text{Si}_2\text{O}_6$	3.067	8.11	0.18	-1.5	0.4	0.224	0.60	1	6	
			4.98	0.13	1.18	0.09	0.456	2.126	2	19	
			7.4	0.4	-0.1	0.2	1.817	2.349	3	6	
			3.7	0.3	1.54	0.07	2.349	4.54	4	22	[8,121,127,231] / [117,165]
Diopside	$\text{CaMgSi}_2\text{O}_6$	3.264	4.9	--	8.4	--	0.201	0.289	1	2	
			7.14	0.03	0.626	0.018	0.289	1.85	2	4	
			6.1	0.2	1.16	0.06	1.85	4.7	4	4	[18,196,213] / [198]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower Up (km/sec)	upper Up (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Beryl	Be <sub>3</sub> Al <sub>2</sub> Si <sub>6</sub> O <sub>18</sub>	2.68	8.83 2.7	0.10 --	0.05 1.6	0.04 --	1.4 4.12	4.12 6.1	2 4	4 2	[187]
<i>Feldspars:</i>											
Orthoclase, Microcline	KAlSi <sub>3</sub> O <sub>8</sub>	2.561	7.70 6.14 3.1	0.11 0.19 0.2	-1.1 0.21 1.39	0.2 0.11 0.05	0.188 1.21 2.41	1.21 2.41 6.27	1 2 4	14 4 4	[12,16,186]
Oligoclase	(NaAlSi <sub>3</sub> -O <sub>8</sub> ) <sub>75</sub> (CaAl <sub>2</sub> -Si <sub>2</sub> O <sub>8</sub> ) <sub>19.5</sub> - (KAlSi <sub>3</sub> -O <sub>8</sub> ) <sub>5.5</sub>	2.635	7.6	0.3	-1.5	1.2	0.195	0.3	1	13	[16]
Anorthite	CaAl <sub>2</sub> Si <sub>2</sub> O <sub>8</sub>	2.769	3.17	0.16	1.45	0.04	2.911	4.338	2	3	[96] / [173,174]
Molten Anorthite <sup>o</sup>	CaAl <sub>2</sub> Si <sub>2</sub> O <sub>8</sub>	2.55	2.85	0.14	1.27	0.09	0.91	2.37	2	6	[168]
Nepheline	(Na,K)AlSiO <sub>4</sub>	2.63	4.76 5.7 2.22	0.08 -- 0.08	0.88 0.1 1.67	0.09 -- 0.02	0.52 1.25 2.57	1.25 1.62 3.94	2 3 4	3 2 4	[187]
<i>Glasses:</i>											
Argutite Glass	GeO <sub>2</sub>	3.655	3.6 0.80	-- 0.06	-0.3 1.79	-- 0.02	0.32 1.35	1.35 4.46	1 2	2 7	[92]
Quartz Glass (fused quartz)	SiO <sub>2</sub>	0.145	1.8 -1.2	-- 0.3	0.36 1.67	-- 0.05	1.789 2.309	2.309 6.507	2 4	2 10	[121]
Quartz Glass	SiO <sub>2</sub>	1.15	0.09	0.2	1.51	0.06	3.07	5.21	4	4	[156]
Quartz Glass	SiO <sub>2</sub>	2.2	0.4	0.5	1.73	0.12	3.61	4.84	4	6	[91]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Molten Diopside <sup>n</sup>	CaMgSi <sub>2</sub> O <sub>6</sub>	2.61	3.30	0.12	1.44	0.08	0.73	2.24	2	5	[168]
Hedenbergite	CaFe <sup>+2</sup> - Si <sub>2</sub> O <sub>6</sub>	3.42	2.8	--	2.8	--	1.1	1.41	2	2	
			6.1	--	0.47	--	1.41	2.42	3	2	
			3.30	0.05	1.620	0.018	2.42	3.62	4	3	[186]
Augite, also Salite	(Ca <sub>0.80</sub> - Na <sub>0.03</sub> )- (Mg <sub>0.76</sub> Fe <sub>0.29</sub> .Ti <sub>0.03</sub> )- (Al <sub>0.20</sub> Si <sub>1.85</sub> -O <sub>6</sub> , CaMg <sub>0.82</sub> - Fe <sub>0.18</sub> Si <sub>2</sub> O <sub>6</sub>	3.435	6.25	0.11	0.85	0.08	0.935	1.87	2	5	
			6.6	0.8	0.6	0.4	1.87	2.48	3	8	
			4.6	1.0	1.4	0.3	2.44	3.82	4	5	[18,196,213]
Jadeite	Na(Al,Fe <sup>+3</sup> )- Si <sub>2</sub> O <sub>6</sub>	3.335	6.41	0.06	1.30	0.08	0.00	1.005	2a	3	
			6.57	0.10	1.09	0.07	0.986	1.94	2b	8	
			7.44	0.12	0.64	0.04	1.94	3.434	4	8	[47,121,135,217]
Spodumene	LiAlSi <sub>2</sub> O <sub>6</sub>	3.14	7.123	0.019	0.007	0.018	0.43	1.45	2	3	
			6.4	--	0.51	--	1.45	2.37	3	2	
			4.0	0.3	1.56	0.09	2.37	3.76	4	3	[187]
Wollastonite	CaSiO <sub>3</sub>	2.82	2.4	0.3	1.65	0.08	1.34	4.06	4	6	[187]
Wollastonite	CaSiO <sub>3</sub>	2.822	6.3	0.2	-0.15	0.15	0.94	1.799	3	5	
			4.1	0.2	1.07	0.09	1.799	2.778	4	4	[121]
Wollastonite	CaSiO <sub>3</sub>	2.89	5.3	--	1.14	--	0.00	1.195	2	2	
			6.826	0.004	-0.156	0.003	1.195	1.933	3	3	
			4.52	0.15	1.03	0.06	1.933	3.282	4	5	[121]
Tremolite	Ca <sub>2</sub> (Mg,- Fe <sup>+2</sup> ) <sub>5</sub> -Si <sub>8</sub> O <sub>22</sub> (OH) <sub>2</sub>	2.901	5.20	0.14	0.91	0.07	1.15	2.82	2	9	
			3.2	0.8	1.6	0.2	2.78	3.94	4	5	[187]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Quartz Glass	SiO <sub>2</sub>	2.204	5.83	0.03	-2.33	0.17	0.00	0.306	1	7	
			5.264	0.019	-0.114	0.014	0.306	2.106	3	32	
			3.40	0.08	0.78	0.03	2.063	2.742	2	20	
			0.80	0.08	1.70	0.02	2.703	5.175	4a	61	
			3.7	0.2	1.196	0.014	5.175	23.64	4b	3	[101,118,121,127, 129,163,217,222] / [104,117,118,175, 177]
Anorthite Glass	CaAl <sub>2</sub> Si <sub>2</sub> O <sub>8</sub>	2.692	6.7	0.3	-0.2	0.4	0.45	1.179	1	5	
			6.43	0.14	0.01	0.07	1.179	2.731	3	3	
			1.8	0.3	1.77	0.07	2.731	4.698	4	15	[52] / [51]
Pyrex (and soda-lime glass)	(SiO <sub>2</sub> ) <sub>81</sub> - (B <sub>2</sub> O <sub>3</sub> ) <sub>12</sub> - (Al <sub>2</sub> O <sub>3</sub> ) <sub>2.5</sub> - (CaO) <sub>0.4</sub> - (MgO) <sub>0.3</sub> - (Na <sub>2</sub> O) <sub>4</sub> - (K <sub>2</sub> O) <sub>1</sub> - (As <sub>2</sub> O <sub>3</sub> ) <sub>0.6</sub> <sup>s</sup>	2.307	5.26	0.14	-0.37	0.15	0.4	1.444	1	5	
			3.96	0.12	0.57	0.06	1.444	2.397	2	4	
			0.6	0.3	1.92	0.09	2.397	4.43	4	11	[95,121,122,136]
Soda-Copper Silicate Glass	(SiO <sub>2</sub> ) <sub>72.2</sub> - (CuO) <sub>12.4</sub> - (Na <sub>2</sub> O) <sub>14</sub> - (Al <sub>2</sub> O <sub>3</sub> ) <sub>0.5</sub> - (SO <sub>3</sub> ) <sub>0.45</sub> - (MgO) <sub>0.1</sub> - (Fe <sub>2</sub> O <sub>3</sub> ) <sub>0.08</sub> <sup>s</sup>	2.48	3.10	0.04	1.28	0.08	0.37	0.71	2	3	
			3.623	0.012	0.531	0.009	0.71	1.55	3	3	
			1.5	0.2	1.95	0.11	1.51	2.57	4	5	[67]
Lunar Glass	(SiO <sub>2</sub> ) <sub>40</sub> - (TiO <sub>2</sub> ) <sub>9</sub> - (Al <sub>2</sub> O <sub>3</sub> ) <sub>11</sub> - (FeO) <sub>17</sub> - (MgO) <sub>10</sub> - (CaO) <sub>11</sub> <sup>s</sup>	1.8	0.10	0.19	1.3	0.2	0.682	1.02	2	4	
			-1.01	0.13	2.39	0.09	0.985	1.825	4	6	[7]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
<b>Other Standards:</b>											
Aluminum: 1100, Russian alloys	Al <sup>r</sup>	1.345	0.00 2.11	0.03 0.19	1.988	0.012	1.323	3.6 6.76	2 4	10 4	[105,142,213]
Aluminum: 1100, Russian alloys	Al <sup>r</sup>	1.885	0.8 3.07	0.4 0.11	2.2 1.39	0.2	1.197	2.77 6.27	2 4	12 6	[39,105,142,213]
Aluminum: 1100, Russian alloys	Al <sup>r</sup>	2.707	6.09 5.44 6.0	0.09 0.04 0.3	-0.1 1.324 1.181	0.4 0.012 0.019	0.01 0.428 5.94	0.526 6.0 30.0	1 2 4	10 59 27	[24,26,27,28,45, 105,115,121,139, 155,162,163,190, 213,228]
Aluminum 2024	Al <sup>r</sup>	1.661	2.20 0.18 2.0	0.12 0.15 0.2	0.19	0.15	0.00	1.012	1	3	
Aluminum 2024	Al <sup>r</sup>	1.955	2.62 0.71 2.8	0.10 0.11 0.2	0.24	0.14	0.00	0.926	1	3	
Aluminum 2024	Al <sup>r</sup>	2.224	3.0 1.3 3.22	-- 0.3 0.16	0.5 2.47 1.58	-- 0.19 0.05	0.00 0.849 1.981	0.849 2.124 4.064	1 2 4	2 7 17	[56,121,134,136]
Aluminum 2024	Al <sup>r</sup>	2.559	4.1 3.24 4.07	-- 0.18 0.10	0.9 2.04 1.54	-- 0.15 0.03	0.00 0.724 1.761	0.724 1.761 3.817	1 2 4	2 5 17	[56,121,134,136]
Aluminum 2024	Al <sup>r</sup>	2.788	5.356	0.011	1.305	0.005	0.00	5.962	2	325	[56,90,121,134,136, 166,213,217,224, 226] / [165]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Molybdenum	Mo	1.277	-0.03	0.06	1.25	0.03	0.678	4.05	2	5	[209]
Molybdenum	Mo	1.72	-0.39	0.10	1.40	0.03	3.02	5.01	2	4	[209]
Molybdenum	Mo	2.55	-0.12	0.05	1.473	0.013	0.654	6.51	2	10	[209]
Molybdenum	Mo	2.914	-0.16	0.04	1.538	0.012	0.65	6.3	2	10	[209]
Molybdenum	Mo	4.435	0.04	0.02	1.735	0.008	0.62	5.66	2	8	[209]
Molybdenum	Mo	5.59	0.30	0.06	1.90	0.02	0.3	5.25	2	13	[209]
Molybdenum	Mo	8.146	0.94 1.91 3.8	0.04 0.18 1.6	2.79 1.98 1.18	0.04 0.09 0.10	0.47 1.13 2.6	1.13 2.6 19.99	2 4a 4b	4 6 3	[162,163,209]
Molybdenum	Mo	10.208	5.14	0.03	1.247	0.005	0.00	20.91	2	61	[24,56,74,108,121, 126,131,136,141, 162,163,166,217, 226]
Molybdenum <sup>m</sup>	Mo	10.208	4.73	0.07	1.45	0.06	0.538	1.414	2	5	[137]
Tantalum	Ta	2.82	0.05 -0.33	0.04 0.03	1.26 1.451	0.03 0.010	0.654 2.06	2.06 4.43	2 4	4 3	[209]
Tantalum	Ta	5.41	-0.25	0.10	1.66	0.04	2.17	3.23	2	3	[209]
Tantalum	Ta	6.20	-0.01	0.02	1.649	0.012	0.601	2.37	2	5	[209]
Tantalum	Ta	8.19	0.028 0.54	0.018 0.09	1.891 1.61	0.012 0.04	1.15 1.8	1.8 3.03	2 4	3 6	[209]
Tantalum	Ta	10.92	0.37 1.43	0.10 0.08	2.23 1.52	0.09 0.04	0.7 1.49	1.49 2.64	2 4	3 5	[209]
Tantalum	Ta	16.649	3.31	0.03	1.306	0.010	0.00	5.86	2	31	[24,86,108,121,136, 139,166,217,226]

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower Up (km/sec)	upper Up (km/sec)	Phase*	No. of Data	References / Temp. Refs.
Tungsten	W	4.6	-0.17	0.14	1.53	0.04	1.45	4.095	2	8	[209]
Tungsten	W	5.5	-0.08	0.07	1.54	0.03	1.38	2.76	2	4	
			0.4	--	1.4	--	2.76	3.63	3	2	
			-3.94	0.19	2.57	0.05	3.63	3.885	4	3	[209]
Tungsten	W	6.57	-0.011	0.013	1.584	0.007	1.295	2.35	2	3	
			1.07	--	1.12	--	2.35	2.68	3	2	
			-1.05	0.14	1.91	0.04	2.68	3.685	4a	4	
			1.7	0.5	1.21	0.04	3.685	16.51	4b	3	[204,209]
Tungsten	W	8.87	0.13	0.05	1.75	0.02	0.56	3.17	2	6	[209]
Tungsten	W	13.36	0.93	0.07	1.97	0.04	0.84	2.54	2	6	[209]
Tungsten	W	18.67	2.86	0.04	2.08	0.06	0.302	0.721	2	4	[121]
Tungsten	W	19.240	4.064	0.010	1.204	0.003	0.00	15.1	2	40	[100,121,131,136, 163,217]
Lexan (polycarbonate)		1.193	1.9	--	2.4	--	0.00	0.421	1	2	
			2.38	0.03	1.551	0.019	0.421	2.53	2	32	
			4.47	0.12	0.70	0.04	2.379	3.651	3	21	
			2.6	0.2	1.27	0.05	3.64	6.92	4	19	[55,101,121]

Notes:

\*Phases: 1) Elastic shock; 2) Low pressure phase; 3) Mixed region; 4) High pressure phase.

a Starting temperature 5K

b Starting temperature 20K

c Starting temperature 75-86K

d Starting temperature 111K

e Starting temperature 122K

f Starting temperature 148K; compressed gas

g Starting temperature 165K

h Starting temperature 196K

**TABLE 1. Shock Wave Equation of State of Minerals and Related Materials of the Solar System**

Mineral	Formula	Sample Density (Mg/m <sup>3</sup> )	C <sub>0</sub> (km/sec)	error ΔC <sub>0</sub> (km/sec)	S	error ΔS	lower U <sub>p</sub> (km/sec)	upper U <sub>p</sub> (km/sec)	Phase*	No. of Data	References / Temp. Refs.
---------	---------	--	----------------------------	--------------------------------------	---	-------------	-------------------------------------	-------------------------------------	--------	-------------	-----------------------------

i Starting temperature 203-230K

j Starting temperature 258-263K

k Starting temperature 273-298K

l Starting temperature 300K; compressed gas

m Starting temperature 1673K

n Starting temperature 1773K

o Starting temperature 1923 K

p 5% cobalt

q raw data not provided

r with impurities

s composition in weight percent oxides

## REFERENCES

- 1 Abe, Y., and T. Matsui, The formation of an impact-generated H<sub>2</sub>O atmosphere and its implications for the early thermal history of the Earth, *Proc. Lunar Planet. Sci. Conf. 15th, Part 2, J. Geophys. Res.*, 90, C545-C559, 1985.
- 2 Ahrens, T. J., High-pressure electrical behavior and equation of state of magnesium oxide from shock wave measurements, *J. Appl. Phys.*, 37, 2532-2541, 1966.
- 3 Ahrens, T. J., Equations of state of iron sulfide and constraints on the sulfur content of the Earth, *J. Geophys. Res.*, 84, 985-998, 1979.
- 4 Ahrens, T. J., Dynamic compression of earth materials, *Science*, 207, 1035-1041, 1980.
- 5 Ahrens, T. J., Application of shock wave data to earth and planetary science, in *Shock Waves in Condensed Matter*, edited by Y. M. Gupta, pp. 571-588, Plenum, New York, 1986.
- 6 Ahrens, T. J., J. D. Bass, and J. R. Abelson, Shock temperatures in metals, in *Shock Compression of Condensed Matter - 1989*, edited by S. C. Schmidt, J. N. Johnson and L. W. Davison, pp. 851-857, Elsevier Publishers, B. V., 1990.
- 7 Ahrens, T. J., and D. M. Cole, Shock compression and adiabatic release of lunar fines from Apollo 17, *Proc. 5th Lunar Science Conf., Suppl. 5, Geochim. et Cosmochim. Acta*, 3, 2333-2345, 1974.
- 8 Ahrens, T. J., and E. S. Gaffney, Dynamic compression of enstatite, *J. Geophys. Res.*, 76, 5504-5513, 1971.
- 9 Ahrens, T. J., W. H. Gust, and E. B. Royce, Material strength effect in the shock compression of alumina, *J. Appl. Phys.*, 39, 4610-4616, 1968.
- 10 Ahrens, T. J., and R. Jeanloz, Pyrite: Shock compression, isentropic release, and composition of the Earth's core, *J. Geophys. Res.*, 92, 10363-10375, 1987.
- 11 Ahrens, T. J., and R. K. Linde, Response of brittle solids to shock compression, in *Behaviour of Dense Media Under High Dynamic Pressures-Symposium on the Behavior of Dense Media under High Dynamic Pressures, Paris, September 1967*, pp. 325-336, Gordon and Breach, New York, 1968.
- 12 Ahrens, T. J., and H.-P. Liu, A shock-induced phase change in orthoclase, *J. Geophys. Res.*, 78, 1274-1278, 1973.
- 13 Ahrens, T. J., and J. D. O'Keefe, Shock melting and vaporization of lunar rocks and minerals, *The Moon*, 4, 214-249, 1972.
- 14 Ahrens, T. J., and J. D. O'Keefe, Equation of state and impact-induced shock-wave attenuation on the Moon, in *Impact and Explosion Cratering*, edited by D. J. Roddy, R. O. Pepin and R. B. Merrill, pp. 639-656, Pergamon, New York, 1977.
- 15 Ahrens, T. J., and C. F. Petersen, Shock wave data and the study of the Earth, in *The Application of Modern Physics to the Earth and Planetary Interiors*, edited by S. K. Runcorn, pp. 449-461, 1969.
- 16 Ahrens, T. J., C. F. Petersen, and J. T. Rosenberg, Shock compression of feldspars, *J. Geophys. Res.*, 74, 2727-2746, 1969.
- 17 Ahrens, T. J., and J. T. Rosenberg, Shock metamorphism: Experiments on quartz and plagioclase, in *Shock Metamorphism of Natural Materials*, edited by B. M. French and N. M. Short, pp. 59-81, Mono Book Corp., Baltimore, 1968.
- 18 Ahrens, T. J., J. T. Rosenberg, and M. H. Ruderman, Stanford Research Institute, Menlo Park, California 94550, DASA 1868, 1966.
- 19 Ahrens, T. J., and M. H. Ruderman, Immersed-foil method for measuring shock wave profiles in solids, *J. Appl. Phys.*, 37, 4758-4765, 1966.
- 20 Ahrens, T. J., H. Tan, and J. D. Bass, Analysis of shock temperature data for iron, *High Pressure Research*, 2, 145-157, 1990.

- 21 Ahrens, T. J., and J. V. G. Gregson, Shock compression of crustal rocks: data for quartz, calcite, and plagioclase rocks, *J. Geophys. Res.*, 69, 4839-4874, 1964.
- 22 Al'tshuler, L. V., Use of shock waves in high-pressure physics, *Sov. Phys. Uspekhi*, 8, 52-91, 1965.
- 23 Al'tshuler, L. V., A. A. Bakanova, and I. P. Dudoladov, Effect of electron structure on the compressibility of metals at high pressure, *Sov. Phys. JETP*, 26, 1115-1120, 1968.
- 24 Al'tshuler, L. V., A. A. Bakanova, I. P. Dudoladov, E. A. Dynin, R. F. Trunin, and B. S. Chekin, Shock adiabatic curves of metals. New data, statistical analysis, and general laws, *J. Appl. Mech. Tech. Phys.*, 2, 145-169, 1981.
- 25 Al'tshuler, L. V., A. A. Bakanova, and R. F. Trunin, Shock adiabats and zero isotherms of seven metals at high pressures, *Sov. Phys. JETP*, 15, 65-74, 1962.
- 26 Al'tshuler, L. V., and S. E. Brusnikin, Equations of state of compressed and heated metals, *High Temp. USSR*, 27, 39-47, 1989.
- 27 Al'tshuler, L. V., N. N. Kalitkin, L. V. Kuz'mina, and B. S. Chekin, Shock adiabats for ultrahigh pressures, *Sov. Phys. JETP*, 45, 167-171, 1977.
- 28 Al'tshuler, L. V., S. B. Kormer, A. A. Bakanova, and R. F. Trunin, Equation of state for aluminum, copper, and lead in the high pressure region, *Sov. Phys. JETP*, 11, 573-579, 1960.
- 29 Al'tshuler, L. V., S. B. Kormer, M. I. Brazhnik, L. A. Vladimirov, M. P. Speranskaya, and A. I. Funtikov, The isentropic compressibility of aluminum, copper, lead, and iron at high pressures, *Sov. Phys. JETP*, 11, 766-775, 1960.
- 30 Al'tshuler, L. V., K. K. Krupnikov, and M. I. Brazhnik, Dynamic compressibility of metals under pressures from 400,000 to 4,000,000 atmospheres, *Sov. Phys. JETP*, 34, 614-619, 1958.
- 31 Al'tshuler, L. V., K. K. Krupnikov, B. N. Ledenev, V. I. Zhuchikhin, and M. I. Brazhnik, Dynamic compressibility and equation of state of iron under high pressure, *Sov. Phys. JETP*, 34, 606-614, 1958.
- 32 Al'tshuler, L. V., L. V. Kuleshova, and M. N. Pavlovskii, The dynamic compressibility, equation of state, and electrical conductivity of sodium chloride at high pressures, *Sov. Phys. JETP*, 12, 10-15, 1961.
- 33 Al'tshuler, L. V., B. N. Moiseev, L. V. Popov, G. V. Simakov, and R. F. Trunin, Relative compressibility of iron and lead at pressures of 31 to 34 Mbar, *Sov. Phys. JETP*, 27, 420-422, 1968.
- 34 Al'tshuler, L. V., M. N. Pavlovskii, and V. P. Draken, Peculiarities of phase transitions in compression and rarefaction shock waves, *Sov. Phys. JETP*, 25, 260-265, 1967.
- 35 Al'tshuler, L. V., M. A. Podurets, G. V. Simakov, and R. F. Trunin, High-density forms of fluorite and rutile, *Soviet Physics--Solid State*, 15, 969-971, 1973.
- 36 Al'tshuler, L. V., G. V. Simakov, and R. F. Trunin, On the chemical composition of the Earth's core, *Izv. Earth Phys.*, 1, 1-3, 1968.
- 37 Al'tshuler, L. V., R. F. Trunin, and G. V. Simakov, Shock-wave compression of periclase and quartz and the composition of the Earth's lower mantle, *Izv. Earth Phys.*, 10, 657-660, 1965.
- 38 Al'tshuler, V. L., M. N. Pavlovskii, L. V. Kuleshova, and G. V. Simakov, Investigation of alkali-metal halides at high pressures and temperatures produced by shock compression, *Soviet Physics--Solid State*, 5, 203-211, 1963.
- 39 Anderson, G. D., D. G. Doran, and A. L. Fahrenbruch, Stanford Research Institute, Menlo Park, California, AFWL-TR-65-147 and AFWL-TR-65-43, 1965.
- 40 Asay, J. R., and M. Shahinpoor (Ed.), *High-Pressure Shock Compression of Solids*, pp. 1-393,

- Springer-Verlag, New York, 1993.
- 41 Bakanova, A. A., V. N. Zubarev, Y. N. Sutulov, and R. F. Trunin, Thermodynamic properties of water at high pressures and temperatures, *Sov. Phys. JETP*, 41, 544-547, 1976.
- 42 Balchan, A. S., and G. R. Cowan, Shock compression of two iron-silicon alloys to 2.7 Megabars, *J. Geophys. Res.*, 71, 3577-3588, 1966.
- 43 Bass, J. D., T. J. Ahrens, J. R. Abelson, and H. Tan, Shock temperature measurements in metals: New results for an Fe alloy, *J. Geophys. Res.*, 95, 21767-76, 1990.
- 44 Bass, J. D., B. Svendsen, and T. J. Ahrens, The temperature of shock-compressed iron, in *High Pressure Research in Mineral Physics*, edited by M. Manghnani and Y. Syono, pp. 393-402, Terra Scientific, Tokyo, 1987.
- 45 Belyakova, M. Y., M. V. Zhernokletov, Y. N. Sutulov, and R. F. Trunin, Shock-wave compression of metal alloys, *Isv. Earth Phys.*, 27, 71-77, 1991.
- 46 Bevington, P. R., *Data reduction and Error Analysis for the Physical Sciences*, pp. 1-336, McGraw-Hill, New York, 1969.
- 47 Birch, F., Compressibility: Elastic Constants, in *Handbook of Physical Constants, revised edition*, edited by S. P. Clark Jr., pp. 153-159, The Geological Society of America, New York, 1966.
- 48 Boness, D. A., Shock-wave experiments and electronic band-structure calculations of materials at high temperature and pressure, Ph.D. thesis, 256 pp., University of Washington, Seattle, 1991.
- 49 Boslough, M., Postshock temperatures in silica, *J. Geophys. Res.*, 93, 6477-6484, 1988.
- 50 Boslough, M. B., T. J. Ahrens, and A. C. Mitchell, Shock temperatures in CaO, *J. Geophys. Res.*, 89, 7845-7851, 1984.
- 51 Boslough, M. B., T. J. Ahrens, and A. C. Mitchell, Shock temperatures in anorthite glass, *Geophys. J.R. astr. Soc.*, 84, 475-489, 1986.
- 52 Boslough, M. B., S. M. Rigden, and T. J. Ahrens, Hugoniot equation of state of anorthite glass and lunar anorthosite, *Geophys. J. R. astr. Soc.*, 84, 455-473, 1986.
- 53 Brown, J. M., T. J. Ahrens, and D. L. Shampine, Hugoniot data for pyrrhotite and the Earth's core, *J. Geophys. Res.*, 89, 6041-6048, 1984.
- 54 Carter, W. J., Hugoniot equation of state of some alkali halides, *High Temp. - High Press.*, 5, 313-318, 1973.
- 55 Carter, W. J., and S. P. Marsh, Hugoniot equations of state of polymers, Los Alamos Scientific Laboratory, unpublished report LA-UR-77-2062, 1977.
- 56 Carter, W. J., S. P. Marsh, J. N. Fritz, and R. G. McQueen, The equation of state of selected materials for high-pressure reference, in *Accurate Characterization of the High Pressure Environment*, edited by E. C. Boyd, pp. 147-158, NBS special pub. 326, Washington D.C., 1971.
- 57 Christian, R. H., R. E. Duff, and F. L. Yarger, Equation of state of gases by shock wave measurements. II. The dissociation energy of nitrogen, *J. Chem. Phys.*, 23, 2045-2049, 1955.
- 58 Coleburn, N. L., Compressibility of pyrolytic graphite, *J. Chem. Phys.*, 40, 71-77, 1964.
- 59 Davison, L., and R. A. Graham, Shock compression of solids, *Phys. Rep.*, 55, 255-379, 1979.
- 60 Dick, R. D., Shock compression of liquid benzene, carbon disulfide, carbon tetrachloride and nitrogen, *Bull. Am. Phys. Soc.*, 13, 579, 1968.
- 61 Dick, R. D., Shock wave compression of benzene, carbon disulfide, carbon tetrachloride and liquid nitrogen, *Los Alamos Scientific Laboratory report, LA-3915*, 1968.
- 62 Dick, R. D., Shock wave compression of benzene, carbon disulfide, carbon tetrachloride, and liquid nitrogen, *J. Chem. Phys.*, 52, 6021-6032, 1970.
- 63 Dick, R. D., and G. I. Kerley, Shock compression data for

- liquids. II. Condensed hydrogen and deuterium, *J. Chem. Phys.*, 73, 5264-5271, 1980.
- 64 Dick, R. D., R. H. Warnes, and J. J. Skalyo, Shock compression of solid argon, *J. Chem. Phys.*, 53, 1648, 1970.
- 65 Doran, D. G., Hugoniot equation of state of pyrolytic graphite to 300 kbars, *J. Appl. Phys.*, 34, 844-851, 1963.
- 66 Doran, D. G., Shock-wave compression of barium titanate and 95/5 lead zirconate titanate, *J. Appl. Phys.*, 39, 40-47, 1968.
- 67 Dremin, A. N., and G. A. Adadurov, The behavior of glass under dynamic loading, *Sov. Phys. - Solid State*, 6, 1379-1384, 1964.
- 68 Duff, R. E., and F. S. Minshall, Investigation of a shock-induced transition in bismuth, *Phys. Rev.*, 108, 1207-1212, 1957.
- 69 Duffy, T. S., T. J. Ahrens, and M. A. Lange, The shock wave equation of state of brucite  $Mg(OH)_2$ , *J. Geophys. Res.*, 96, 14,319-14,330, 1991.
- 70 Duvall, G. E., and G. R. Fowles, Shock waves, in *High Pressure Physics and Chemistry*, vol. 2, edited by R. S. Bradley, pp. 209-292, Academic Press, New York, 1963.
- 71 Fortov, V. E., A. A. Leont'ev, A. N. Dremin, and V. K. Gryaznov, Shock-wave production of a non-ideal plasma, *Sov. Phys. JETP*, 44, 116-122, 1976.
- 72 Fowles, R., Dynamic compression of quartz, *J. Geophys. Res.*, 72, 5729-5742, 1967.
- 73 Fritz, J. N., S. P. Marsh, W. J. Carter, and R. G. McQueen, The Hugoniot equation of state of sodium chloride in the sodium chloride structure in, in *Accurate Characterization of the High-Pressure Environment, proceedings of a symposium held at the National Bureau of Standards, Gaithersburg, Maryland, Special publication*, vol. 326, NBS, Washington DC, 1971.
- 74 Furnish, M. D., L. C. Chhabildas, D. J. Steinberg, and G. T. G. III, Dynamic behavior of fully dense molybdenum, in *Shock Compression of Condensed Matter 1991*, edited by S. C. Schmidt, R. D. Dick, J. W. Forbes and D. G. Tasker, pp. 419-422, Elsevier Science Publishers B.V, 1992.
- 75 Gaffney, E. S., and T. J. Ahrens, Identification of ice VI on the Hugoniot of ice I<sub>h</sub>, *Geophys. Res. Lett.*, 7, 407-409, 1980.
- 76 Goto, T., T. J. Ahrens, G. R. Rossman, and Y. Syono, Absorption spectrum of shock-compressed  $Fe^{2+}$ -bearing MgO and the radiative conductivity of the lower mantle, *Phys. Earth Planet. Inter.*, 22, 277-288, 1980.
- 77 Graham, E. K., and T. J. Ahrens, Shock wave compression of iron-silicate garnet, *J. Geophys. Res.*, 78, 375-392, 1973.
- 78 Graham, R. A., and W. P. Brooks, Shock-wave compression of sapphire from 15 to 420 kbar. The effects of large anisotropic compressions, *J. Phys. Chem Solids*, 32, 2311-2330, 1971.
- 79 Graham, R. A., O. E. Jones, and J. R. Holland, Physical behavior of germanium under shock wave compression, *J. Phys. Chem. Solids*, 27, 1519-1529, 1966.
- 80 Grigor'ev, F. V., S. B. Kormer, O. L. Mikhailova, M. A. Mochalov, and V. D. Urlin, Shock compression and brightness temperature of a shock wave front in argon. Electron screening of radiation, *Sov. Phys. JETP*, 61, 751-757, 1985.
- 81 Gryaznov, V. K., M. V. Zhernokletov, V. N. Zubarev, I. L. Ioselevskii, and V. E. Fortov, Thermodynamic properties of a nonideal argon or xenon plasma, *Sov. Phys. JETP*, 51, 288-295, 1980.
- 82 Gust, W. H., Phase transition and shock-compression parameters to 120 GPa for three types of graphite and for amorphous carbon, *Phys. Rev. B*, 22, 4744-4756, 1980.
- 83 Harvey, W. B., and R. D. Dick, Shock compression of liquid ammonia, *Bull. Am. Phys. Soc.*, 20, 48, 1975.
- 84 Hauver, G. E., and A. Melani, Ballistics Research Labs,

- Aberdeen Proving Ground, Maryland, BRL MR 2061, 1970.
- 85 Hayes, D. B., Polymorphic phase transformation rates in shock-loaded potassium chloride, *J. Appl. Phys.*, **45**, 1208-1217, 1974.
- 86 Holmes, N. C., J. A. Moriarty, G. R. Gathers, and W. J. Nellis, The equation of state of platinum to 660 GPa (6.6 Mbar), *J. Appl. Phys.*, **66**, 2962-2967, 1989.
- 87 Holmes, N. C., W. J. Nellis, W. B. Graham, and G. E. Walrafen, Spontaneous Raman spectroscopy of shocked H<sub>2</sub>O, *Physica*, **139 and 140B**, 568-570, 1986.
- 88 Holmes, N. C., and E. F. See, Shock compression of low-density microcellular materials, in *Shock Compression of Condensed Matter 1991*, edited by S. C. Schmidt, R. D. Dick, J. W. Forbes and D. G. Tasker, pp. 91-94, Elsevier Science Publishers B.V., 1992.
- 89 Hughes, D. S., L. E. Gourley, and M. F. Gourley, Shock-wave compression of iron and bismuth, *J. Appl. Phys.*, **32**, 624-629, 1961.
- 90 Isbell, W. M., F. H. Shipman, and A. H. Jones, Hugoniot equation of state measurements for eleven materials to five Megabars, *DASA File No 15.025, MSL-68-13*, 128 pp., 1968.
- 91 Jackson, I., and T. J. Ahrens, Shock-wave compression of single crystal forsterite, *J. Geophys. Res.*, **84**, 3039-3048, 1979.
- 92 Jackson, I., and T. J. Ahrens, Shock-wave compression of vitreous and rutile-type GeO<sub>2</sub>: A comparative study, *Phys. Earth Planet. Int.*, **20**, 60-70, 1979.
- 93 Jeanloz, R., Shock wave equation of state and finite strain theory, *J. Geophys. Res.*, **94**, 5873-5886, 1989.
- 94 Jeanloz, R., The nature of the Earth's core, *Ann. Rev. Earth and Planet. Sci.*, **18**, 357-386, 1990.
- 95 Jeanloz, R., and T. J. Ahrens, Pyroxenes and olivines: Structural implications of shock-wave data for high pressure phases, in *High-Pressure Research: Applications to Geophysics*, edited by M. H. Manghnani and S. Akimoto, pp. 439-462, Academic, San Diego, Calif., 1977.
- 96 Jeanloz, R., and T. J. Ahrens, Anorthite, thermal equation of state to high pressures, *Geophys. J. R. astr. Soc.*, **62**, 529-549, 1980.
- 97 Jeanloz, R., and T. J. Ahrens, Equations of state of FeO and CaO, *Geophys. J. R. astron. Soc.*, **62**, 505-528, 1980.
- 98 Jeanloz, R., and R. Grover, Birch-Murnaghan and Us-Up equations of state, in *Proceedings of the American Physical Society Topical Conference on Shock Waves in Condensed Matter, Monterey, CA, 1987*, edited by S. C. Schmidt and N. C. Holmes, pp. 69-72, Plenum, New York, 1988.
- 99 Johnson, J. O., and J. Wackerle, Shock-wave compression of porous magnesium and ammonium sulfate, in *Behaviour of Dense Media Under High Dynamic Pressures-Symposium on the Behavior of Dense Media under High Dynamic Pressures, Paris, September 1967*, pp. 217-226, Gordon and Breach, New York, 1968.
- 100 Jones, A. H., W. M. Isbell, and C. J. Maiden, Measurement of the very-high-pressure properties of materials using a light-gas gun, *J. Appl. Phys.*, **37**, 3493-3499, 1966.
- 101 Jones, A. H., W. M. Isbell, F. H. Shipman, R. D. Perkins, S. J. Green, and C. J. Maiden, Material properties measurements for selected materials, NASA Ames, Interim Report, NAS2-3427, MSL-68-9, 55 pp., 1968.
- 102 Kalashnikov, N. G., M. N. Pavlovskiy, G. V. Simakov, and R. F. Trunin, Dynamic compressibility of calcite-group minerals, *Izv. Earth Phys.*, **2**, 23-29, 1973.
- 103 Kondo, K.-I., and T. J. Ahrens, Heterogeneous shock-induced thermal radiation in minerals, *Phys. Chem. Minerals*, **9**, 173-181, 1983.
- 104 Kondo, K.-I., T. J. Ahrens, and A. Sawaoka, Shock-induced radiation spectra of fused quartz,

- J. Appl. Phys.*, 54, 4382-4385, 1983.
- 105 Kormer, S. B., A. I. Funtikov, V. D. Urlin, and A. N. Kolesnikova, Dynamic compression of porous metals and the equation of state with variable specific heat at high temperatures, *Sov. Phys. JETP*, 15, 477-488, 1962.
- 106 Kormer, S. B., M. V. Sinitsyn, A. I. Funtikov, V. D. Urlin, and A. V. Blinov, Investigation of the compressibility of five ionic compounds at pressures up to 5 Mbar, *Sov. Phys. JETP*, 20, 811-819, 1965.
- 107 Kormer, S. B., M. V. Sinitsyn, G. A. Kirilov, and V. D. Urlin, Experimental determination of temperature in shock-compressed NaCl and KCl and of their melting curves at pressures up to 700 kbar, *Soviet Phys. JETP*, 21, 689-708, 1965.
- 108 Krupnikov, K. K., A. A. Bakanova, M. I. Brazhnik, and R. F. Trunin, An investigation of the shock compressibility of titanium, molybdenum, tantalum, and iron, *Sov. Phys. Doklady*, 2, 205-208, 1963.
- 109 Lagus, P. L., and T. J. Ahrens, Shock wave measurements on solid hydrogen and argon, *J. Chem. Phys.*, 59, 3517-3522, 1973.
- 110 Lange, M., and T. J. Ahrens, Fragmentation of ice by low velocity impact, in *12th Lunar and Planetary Science Conference*, pp. 1667-1687, Pergamon Press, Houston, TX, 1981.
- 111 Lange, M. A., and T. J. Ahrens, FeO and H<sub>2</sub>O and the homogeneous accretion of the Earth, *Earth Planet. Sci. Lett.*, 71, 111-119, 1984.
- 112 Lange, M. A., P. Lambert, and T. J. Ahrens, Shock effects on hydrous minerals and implications for carbonaceous meteorites, *Geochim. Cosmochim. Acta*, 49, 1715-1726, 1985.
- 113 Larson, D. B., A shock-induced phase transformation in bismuth, *J. Appl. Phys.*, 38, 1541-1546, 1967.
- 114 Larson, D. B., G. D. Bearson, and J. R. Taylor, Shock-wave studies of ice and two frozen soils, in *North American Contribution to the Second International Conference on Permafrost*, edited by T. L. Péwé and J. R. Mackay, pp. 318-325, 1973.
- 115 Lundberg, C. D., and W. Herrmann, Equation of state of 6061-T6 aluminum at low pressures, *J. Appl. Phys.*, 34, 2046-2052, 1963.
- 116 Lysne, P. C., A comparison of calculated and measured low-stress Hugoniots and release adiabats of dry and water-saturated tuff, *J. Geophys. Res.*, 75, 4375-4386, 1970.
- 117 Lyzenga, G. A., and T. J. Ahrens, Shock temperature measurements in Mg<sub>2</sub>SiO<sub>4</sub> and SiO<sub>2</sub> at high pressures, *Geophys. Res. Lett.*, 7, 141-144, 1980.
- 118 Lyzenga, G. A., T. J. Ahrens, and A. C. Mitchell, Shock temperatures of SiO<sub>2</sub> and their geophysical implications, *J. Geophys. Res.*, 88, 2431-2444, 1983.
- 119 Lyzenga, G. A., T. J. Ahrens, W. J. Nellis, and A. C. Mitchell, The temperature of shock-compressed water, *J. Chem. Phys.*, 76, 6282-6286, 1982.
- 120 Marsh, S. P., Hugoniot equation of state of beryllium oxide, *High Temp.-High Press.*, 5, 503-508, 1973.
- 121 Marsh, S. P. (Ed.), *LASL Shock Hugoniot Data*, pp. 1-658, University of California Press, Berkeley, 1980.
- 122 Mashimo, T., Y. Hanaoka, and K. Nagayama, Elastoplastic properties under shock compression of Al<sub>2</sub>O<sub>3</sub> single crystal and polycrystal, *J. Appl. Phys.*, 63, 327-336, 1988.
- 123 Mashimo, T., K. Nagayama, and A. Sawaoka, Anisotropic elastic limits and phase transitions of rutile phase TiO<sub>2</sub> under shock compression, *J. Appl. Phys.*, 54, 5043-5048, 1983.
- 124 Mashimo, T., K. Nagayama, and A. Sawaoka, Shock

- compression of zirconia  $ZrO_2$  and zircon  $ZrSiO_4$  in the pressure range up to 150 GPa, *Phys. Chem. Minerals*, 9, 237-247, 1983.
- 125 McMahan, A. K., B. L. Hord, and M. Ross, Experimental and theoretical study of metallic iodine, *Phys. Rev. B* 15, 726-737, 1977.
- 126 McQueen, R. G., Laboratory techniques for very high pressures and the behavior of metals under dynamic loading in, in *Metallurgy at High Pressures and High Temperatures*, edited by J. K. A. Gschneider, M. T. Hepworth and N. A. D. Parlee, pp. 44-132, Gordon and Breach, New York, 1964.
- 127 McQueen, R. G., The equation of state of mixtures, alloys, and compounds, in *Seismic Coupling- Proceedings of a meeting sponsored by the Advanced Research Projects Agency, January 15-16, 1968*, edited by G. Simmons, Stanford Research Institute, Menlo Park, CA, 1968.
- 128 McQueen, R. G., W. J. Carter, J. N. Fritz, and S. P. Marsh, The solid-liquid phase line in Cu, in *Accurate Characterization of the High-Pressure Environment*, proceedings of a symposium held at the National Bureau of Standards, Gaithersburg, Maryland, October 14-18, 1968,
- 129 Special publication 326, NBS, Washington DC, 1971.
- 130 McQueen, R. G., J. C. Jamieson, and S. P. Marsh, Shock-wave compression and X-ray studies of titanium dioxide, *Science*, 155, 1401-1404, 1967.
- 131 McQueen, R. G., and S. P. Marsh, Equation of state for nineteen metallic elements from shock-wave measurements to 2 Megabars, *J. Appl. Phys.*, 31, 1253-1269, 1960.
- 132 McQueen, R. G., and S. P. Marsh, Shock-wave compression of iron-nickel alloys and the Earth's core, *J. Geophys. Res.*, 71, 1751-1756, 1966.
- 133 McQueen, R. G., and S. P. Marsh, Hugoniot of graphite of various initial densities and the equation of state of carbon, in *Behavior of Dense Media under High Dynamic Pressures, Symposium H. D. P.*, September 1967, Gordon and Breach, New York, Paris, 1968.
- 134 McQueen, R. G., S. P. Marsh, and W. J. Carter, The determination of new standards for shock wave equation-of-state work, in *Behavior of Dense Media under High Dynamic Pressures, Symposium on the Behavior of Dense Media under High Dynamic Pressures, Paris, September 1967*, pp. 67-83,
- Gordon and Breach, New York, 1968.
- 135 McQueen, R. G., S. P. Marsh, and J. N. Fritz, Hugoniot equation of state of twelve rocks, *J. Geophys. Res.*, 72, 4999-5036, 1967.
- 136 McQueen, R. G., S. P. Marsh, J. W. Taylor, J. N. Fritz, and W. J. Carter, The equation of state of solids from shock wave studies, in *High-Velocity Impact Phenomena*, edited by R. Kinslow, pp. 249-419, Academic Press, New York, 1970.
- 137 Miller, G. H., T. J. Ahrens, and E. M. Stolper, The equation of state of molybdenum at 1400°C, *J. Appl. Phys.*, 63, 4469-4475, 1988.
- 138 Minshall, F. S., The dynamic response of iron and iron alloys to shock waves, *Met. Soc. Conf.*, 9, 249-274, 1961.
- 139 Mitchell, A. C., and W. J. Nellis, Shock compression of aluminum, copper and tantalum, *J. Appl. Phys.*, 52, 3363-3374, 1981.
- 140 Mitchell, A. C., and W. J. Nellis, Equation of state and electrical conductivity of water and ammonia shocked to the 100 GPa (1 Mbar) pressure range, *J. Chem. Phys.*, 76, 6273-6281, 1982.
- 141 Mitchell, A. C., W. J. Nellis, J. A. Moriarty, R. A. Heinle, N. C. Holmes, R. E. Tipton, and G. W. Repp, Equation of state of Al, Cu, Mo, and Pb at shock

- pressures up to 2.4 TPa (24 Mbar), *J. Appl. Phys.*, **69**, 2981-2986, 1991.
- 142 Morgan, D. T., M. Rockowitz, and A. L. Atkinson, Avco Corporation, Research and Development Division, Wilmington, Massachusetts, AFWL-TR-65-117, 1965.
- 143 Munson, D. E., and L. M. Barker, Dynamically determined pressure-volume relationships for aluminum, copper, and lead, *J. Appl. Phys.*, **37**, 1652-1660, 1966.
- 144 Nellis, N. J., A. C. Mitchell, M. v. Thiel, G. J. Devine, and R. J. Trainor, Equation-of-state data for molecular hydrogen and deuterium at shock pressures in the range 2-76 GPa (20-760 kbar), *J. Chem. Phys.*, **79**, 1480-1486, 1983.
- 145 Nellis, W. J., N. C. Holmes, A. C. Mitchell, R. J. Trainor, G. K. Governo, M. Ross, and D. A. Young, Shock compression of liquid helium to 56 GPa (560 kbar), *Phys. Rev. Lett.*, **53**, 1248-1251, 1953.
- 146 Nellis, W. J., and A. C. Mitchell, Shock compression of liquid argon, nitrogen, and oxygen to 90 GPa (900 kbar), *J. Chem. Phys.*, **73**, 6137-6145, 1980.
- 147 Nellis, W. J., A. C. Mitchell, F. H. Ree, M. Ross, N. C. Holmes, R. J. Trainor, and D. J. Erskine, Equation of state of shock-compressed liquids: carbon dioxide and air, *J. Chem. Phys.*, **95**, 5268-5272, 1991.
- 148 Nellis, W. J., J. A. Moriarty, A. C. Mitchell, M. Ross, R. G. Dandrea, N. W. Ashcroft, N. C. Holmes, and G. R. Gathers, Metals physics at ultrahigh pressures: aluminum, copper and lead as prototypes, *Phys. Rev. Lett.*, **60**, 1414-1417, 1988.
- 149 Nellis, W. J., H. B. Radousky, D. C. Hamilton, A. C. Mitchell, N. C. Holmes, K. B. Christianson, and M. van Thiel, Equation of state, shock temperature, and electrical-conductivity data of dense fluid nitrogen in the region of the dissociative phase transition, *J. Chem. Phys.*, **94**, 2244-2257, 1991.
- 150 Nellis, W. J., F. H. Ree, M. van Thiel, and A. C. Mitchell, Shock compression of liquid carbon monoxide and methane to 90 GPa (900 kbar), *J. Chem. Phys.*, **75**, 3055-3063, 1981.
- 151 Nellis, W. J., M. v. Thiel, and A. C. Mitchell, Shock compression of liquid xenon to 130 GPa (1.3 Mbar), *Phys. Rev. Lett.*, **48**, 816-818, 1982.
- 152 Pavlovskii, M. N., Shock compressibility of six very hard substances, *Soviet Physics--Solid State*, **12**, 1736-1737, 1971.
- 153 Pavlovskii, M. N., Shock compression of diamond, *Sov. Phys. Solid State*, **13**, 741-742, 1971.
- 154 Pavlovskii, M. N., V. Y. Vashchenko, and G. V. Simakov, Equation of state for cesium iodide, *Sov. Phys. - Solid State*, **7**, 972-974, 1965.
- 155 Podurets, M. A., L. V. Popov, A. G. Sevast'yanova, G. V. Simakov, and R. F. Trunin, On the relation between the size of studied specimens and the position of the silica shock adiabat, *Izv. Earth Phys.*, **11**, 727-728, 1976.
- 156 Podurets, M. A., G. V. Simakov, G. S. Telegin, and R. F. Trunin, Polymorphism of silica in shock waves and equation of state of coesite and stishovite, *Izv. Earth Phys.*, **17**, 9-15, 1981.
- 157 Podurets, M. A., G. V. Simakov, R. F. Trunin, L. V. Popov, and B. N. Moiseev, Compression of water by strong shock waves, *Sov. Phys. JETP*, **35**, 375-376, 1972.
- 158 Rabie, R., and J. J. Dick, Equation of state and crushing dynamics of low-density silica aerogels, in *Shock Compression of Condensed Matter 1991*, edited by S. C. Schmidt, R. D. Dick, J. W. Forbes and D. G. Tasker, pp. 87-90, Elsevier Science Publishers B.V., 1992.
- 159 Radousky, H. B., A. C. Mitchell, and W. J. Nellis, Shock temperature measurements of planetary ices: NH<sub>3</sub>, CH<sub>4</sub>, and "Synthetic Uranus", *Chem. Phys.*, **93**, 8235-8239, 1990.

- 160 Radousky, H. B., W. J. Nellis, M. Ross, D. C. Hamilton, and A. C. Mitchell, Molecular dissociation and shock-induced cooling in fluid nitrogen at high densities and temperatures, *Phys. Rev. Lett.*, 57, 2419-2422, 1986.
- 161 Radousky, H. B., and M. Ross, Shock temperature measurements in high density fluid xenon, *Phys. Lett. A*, 129, 43-46, 1988.
- 162 Ragan, C. E., III, Shock-wave experiments at threefold compression, *Phys. Rev. A*, 29, 1391-1402, 1984.
- 163 Ragan, C. E. I., Shock compression measurements at 1 to 7 TPa, *Phys. Rev. A*, 82, 3360-3375, 1982.
- 164 Raikes, S. A., and T. J. Ahrens, Measurements of post-shock temperatures in aluminum and stainless steel, in *High Pressure Science and Technology*, edited by K. D. Timmerhaus and M. S. Barber, pp. 889-894, Plenum Press, New York, 1979.
- 165 Raikes, S. A., and T. J. Ahrens, Post-shock temperature in minerals, *Geophys. J. Roy. Astr. Soc.*, 58, 717-747, 1979.
- 166 Rice, M. H., R. G. McQueen, and J. M. Walsh, Compression of solids by strong shock waves, in *Solid State Phys.*, vol. 6, edited by F. Seitz and D. Turnbull, pp. 1-63, Academic Press, New York, 1958.
- 167 Rice, M. H., and J. M. Walsh, Equation of state of water to 250 kilobars, *J. Chem. Phys.*, 26, 824-830, 1957.
- 168 Rigden, S. M., T. J. Ahrens, and E. M. Stolper, High-pressure equation of state of molten anorthite and diopside, *J. Geophys. Res.*, 94, 9508-9522, 1989.
- 169 Ringwood, A. E., *Origin of the Earth and Moon*, pp. 1-295, Springer-Verlag, Berlin, New York, 1979.
- 170 Roddy, D. J., R. O. Pepin, and R. B. Merrill, *Impact and Explosion Cratering*, pp. 1-1301, Pergamon, New York, 1977.
- 171 Ruoff, A. L., Linear shock-velocity-particle-velocity relationship, *J. Appl. Phys.*, 38, 4976-4980, 1967.
- 172 Schmidt, D. N., and R. K. Linde, Response of distended copper, iron, and tungsten to shock loading, SRI, Air Force Weapons Lab Report, AF 29(601)-7236, 1968.
- 173 Schmitt, D. R., and T. J. Ahrens, Temperatures of shock-induced shear instabilities and their relationship to fusion curves, *Geophys. Res. Lett.*, 10, 1077-1080, 1983.
- 174 Schmitt, D. R., and T. J. Ahrens, Emission spectra of shock compressed solids, in *Shock Waves in Condensed Matter-1983*, vol. Chapter VII:6, edited by J. R. Asay, R. A. Graham and G. K. Straub, pp. 513-516, Elsevier Science Publishers B.V., 1984.
- 175 Schmitt, D. R., and T. J. Ahrens, Shock temperatures in silica glass: Implications for modes of shock-induced deformation, phase transformation, and melting with pressure, *J. Geophys. Res.*, 94, 5851-5871, 1989.
- 176 Schmitt, D. R., T. J. Ahrens, and B. Svendsen, Shock-induced melting and shear banding in single crystal NaCl, *J. Appl. Phys.*, 63, 99-106, 1988.
- 177 Schmitt, D. R., B. Svendsen, and T. J. Ahrens, Shock-induced radiation from minerals, in *Shock Waves in Condensed Matter*, edited by Y. M. Gupta, pp. 261-265, Plenum Press, New York, 1986.
- 178 Schott, G. L., Shock-compressed carbon dioxide: liquid measurements and comparisons with selected models, *High Pressure Res.*, 6, 187-200, 1991.
- 179 Schott, G. L., M. S. Shaw, and J. D. Johnson, Shocked states from initially liquid oxygen-nitrogen systems, *J. Chem. Phys.*, 82, [4264-4275], 1982.
- 180 Seitz, W. L., and J. Wackerle, Reflected-shock Hugoniot for liquid argon between 0.26 and 0.74 Megabars, *Bull. Am. Phys. Soc.*, 17, 1093, 1972.
- 181 Sekine, T., and T. J. Ahrens, Equation of state of heated glassy

- carbon, in, in *Shock Compression of Condensed Matter 1991*, edited by R. D. D. S. C. Schmidt J. W. Forbes, and D. G. Tasker, pp. 57-60, 1992.
- 182 Sekine, T., A. M. Rubin, and T. J. Ahrens, Shock wave equation of state of muscovite, *J. Geophys. Res.*, 96, 19675-19680, 1991.
- 183 Sharpton, V. L., and P. D. Ward (Ed.), *Global Catastrophes in Earth History: An Interdisciplinary Conference on Impacts, Volcanism, and Mass Mortality*, pp. 1-631, The Geological Society of America, Inc., Special Paper 247, Boulder, Colorado, 1990.
- 184 Shipman, F. H., W. M. Isbell, and A. H. Jones, High pressure Hugoniot measurements for several Nevada test site rocks, DASA Report 2214, MSL-68-15, 114 pp., 1969.
- 185 Silver, L. T., and P. Schultz (Ed.), *Geological Implications of Impacts of Large Asteroids and Comets on the Earth*, pp. 1-528, The Geological Society of America, Special Paper 190, Boulder, Colorado, 1982.
- 186 Simakov, G. V., M. N. Pavlovskiy, N. G. Kalashnikov, and R. F. Trunin, Shock compressibility of twelve minerals, *Izv. Earth Phys.*, 10, 488-492, 1974.
- 187 Simakov, G. V., and R. F. Trunin, Compression of minerals by shock waves, *Izv. Earth Phys.*, 16, 134-137, 1980.
- 188 Simakov, G. V., and R. F. Trunin, Shockwave compression of ultraporous silica, *Izv. Earth Physics*, 26, 952-956, 1990.
- 189 Skalyo, J., Jr., R. D. Dick, and R. H. Warnes, Shock compression of solid argon, *Bull. Am. Phys. Soc.*, 13, 579, 1968.
- 190 Skidmore, I. C., and E. Morris, in *Thermodynamics of Nuclear Materials*, pp. 173, International Atomic Energy Agency, Vienna, Austria, 1962.
- 191 Somerville, M., and T. J. Ahrens, Shock compression of  $KFeS_2$  and the question of potassium in the core, *J. Geophys. Res.*, 85, 7016-7024, 1980.
- 192 Stevenson, D. J., Models of the earth's core, *Science*, 214, 611-619, 1981.
- 193 Stöffler, D., Deformation and transformation of rock-forming minerals by natural and experimental shock processes, I., *Fortschr. Miner.*, 49, 50-113, 1972.
- 194 Stöffler, D., Deformation and transformation of rock-forming minerals by natural and experimental shock processes. II. Physical properties of shocked minerals, *Fortschr. Miner.*, 51, 256-289, 1974.
- 195 Stolper, E. M., D. Walker, B. H. Hager, and J. F. Hays, Melt segregation from partially molten source regions: The importance of melt density and source region size, *J. Geophys. Res.*, 86, 6261-6271, 1981.
- 196 Svendsen, B., and T. J. Ahrens, Dynamic compression of diopside and salite to 200 GPa, *Geophys. Res. Lett.*, 10, 501-504, 1983.
- 197 Svendsen, B., and T. J. Ahrens, Shock-induced temperatures of MgO, *Geophys. J. R. Astr. Soc.*, 91, 667-691, 1987.
- 198 Svendsen, B., and T. J. Ahrens, Shock-induced temperatures of  $CaMgSi_2O_6$ , *J. Geophys. Res.*, 95, 6943-6953, 1990.
- 199 Swenson, C. A., J. W. Shaner, and J. M. Brown, Hugoniot overtake sound-velocity measurements on CsI, *Phys. Rev. B*, 34, 7924-7935, 1986.
- 200 Syono, Y., T. Goto, and Y. Nakagawa, Phase-transition pressures of  $Fe_3O_4$  and GaAs determined from shock-compression experiments, in *High Pressure Research: Application in Geophysics*, edited by M. H. Manghnani and S. Akimoto, pp. 463-476, Academic Press, New York, 1977.
- 201 Syono, Y., T. Goto, J.-I. Sato, and H. Takei, Shock compression measurements of single-crystal forsterite in the pressure range 15-93 GPa, *J. Geophys. Res.*, 86, 6181-6186, 1981.

- 202 Taylor, J. W., and M. H. Rice, Elastic-plastic properties of iron, *J. Appl. Phys.*, 34, 364-371, 1963.
- 203 Trunin, R. F., Compressibility of various substances at high shock pressures: an overview, *Izv. Earth Phys.*, 22, 103-114, 1986.
- 204 Trunin, R. F., A. B. Medvedev, A. I. Funtikov, M. A. Podurets, G. V. Simakov, and A. G. Sevast'yanov, Shock compression of porous iron, copper, and tungsten, and their equation of state in the terapascal pressure range, *Sov. Phys. JETP*, 68, 356-361, 1989.
- 205 Trunin, R. F., M. A. Podurets, B. N. Moiseev, G. V. Simakov, and L. V. Popov, Relative compressibility of copper, cadmium, and lead at high pressures, *Sov. Phys. JETP*, 29, 630-631, 1969.
- 206 Trunin, R. F., M. A. Podurets, G. V. Simakov, L. V. Popov, and B. N. Moiseev, An experimental verification of the Thomas-Fermi model for metals under high pressure, *Sov. Phys. JETP*, 35, 550-552, 1972.
- 207 Trunin, R. F., G. V. Simakov, and M. A. Podurets, Compression of porous quartz by strong shock waves, *Izv. Earth Phys.*, 2, 102-106, 1971.
- 208 Trunin, R. F., G. V. Simakov, M. A. Podurets, B. N. Moiseyev, and L. V. Popov, Dynamic compressibility of quartz and quartzite at high pressure, *Izv. Earth Phys.*, 1, 8-12, 1971.
- 209 Trunin, R. F., G. V. Simakov, Y. N. Sutulov, A. B. Medvedev, B. D. Rogozkin, and Y. E. Fedorov, Compressibility of porous metals in shock waves, *Sov. Phys. JETP*, 69, 580-588, 1989.
- 210 Trunin, R. F., M. V. Zhernokleto, N. F. Kuznetsov, G. V. Simakov, and V. V. Shutov, Dynamic compressibility of aqueous solutions of salts, *Izv. Earth Physics*, 23, 992-995, 1987.
- 211 Tyburczy, J. A., T. S. Duffy, T. J. Ahrens, and M. A. Lange, Shock wave equation of state of serpentine to 150 GPa: Implications of the occurrence of water in the Earth's lower mantle, *J. Geophys. Res.*, 96, 18011-18027, 1991.
- 212 Urlin, V. D., M. A. Machalov, and O. L. Mikhailova, Liquid xenon study under shock and quasi-isentropic compression, *High Pressure Research*, 8, 595-605, 1992.
- 213 van Thiel, M., (editor), Compendium of Shock Wave Data, University of California, Lawrence Livermore Laboratory, UCRL-50801, Vol. 1, Rev. 1, 755 pp., 1977.
- 214 Van Thiel, M., and B. J. Alder, Shock compression of argon, *J. Chem. Phys.*, 44, 1056-1065, 1966.
- 215 van Thiel, M., and B. J. Alder, Shock compression of liquid hydrogen, *Mol. Phys.*, 19, 427-435, 1966.
- 216 Van Thiel, M., L. B. Hord, W. H. Gust, A. C. Mitchell, M. D'Addario, B. K. E. Wilbarger, and B. Barrett, Shock compression of deuterium to 900 kbar, *Phys. Earth Planet. Int.*, 9, 57-77, 1974.
- 217 van Thiel, M., A. S. Kusubov, and A. C. Mitchell, Compendium of Shock Wave Data, Lawrence Radiation Laboratory (Livermore), UCRL-50108, 1967.
- 218 Vassiliou, M. S., and T. J. Ahrens, Hugoniot equation of state of periclase to 200 GPa, *Geophys. Res. Lett.*, 8, 729-732, 1981.
- 219 Vassiliou, M. S., and T. J. Ahrens, The equation of state of  $Mg_{0.6}Fe_{0.4}O$  to 200 GPa, *Geophys. Res. Lett.*, 9, 127-130, 1982.
- 220 Vizgirda, J., and T. J. Ahrens, Shock compression of aragonite and implications for the equation of state of carbonates, *J. Geophys. Res.*, 87, 4747-4758, 1982.
- 221 Voskoboinikov, I. M., M. F. Gogulya, and Y. A. Dolgorobodov, Temperatures of shock compression of liquid nitrogen and argon, *Sov. Phys. Doklady*, 24, 375-376, 1980.

- 222 Wackerle, J., Shock-wave compression of quartz, *J. Appl. Phys.*, 33, 922-937, 1962.
- 223 Wackerle, J., W. L. Seitz, and J. C. Jamieson, Shock-wave equation of state for high-density oxygen in, in *Behavior of Dense Media under High Dynamic Pressures Symposium, Paris, September 1967*, pp. 85, Gordon and Breach, New York, 1968.
- 224 Walsh, J. M., and R. H. Christian, Equation of state of metals from shock wave measurements, *Phys. Rev.*, 97, 1544-1556, 1955.
- 225 Walsh, J. M., and M. H. Rice, Dynamic compression of liquids from measurements of strong shock waves, *J. Chem. Phys.*, 26, 815-823, 1957.
- 226 Walsh, J. M., M. H. Rice, R. G. McQueen, and F. L. Yarger, Shock-wave compressions of twenty-seven metals. Equation of state of metals., *Phys. Rev.*, 108, 196-216, 1957.
- 227 Warnes, R. H., Investigation of a shock-induced phase transition in antimony, *J. Appl. Phys.*, 38, 4629-4631, 1967.
- 228 Warnes, R. H., Shock wave compression of three polynuclear aromatic compounds, *J. Chem. Phys.*, 53, 1088, 1970.
- 229 Watt, J. P., and T. J. Ahrens, Shock compression of single-crystal forsterite, *J. Geophys. Res.*, 88, 9500-9512, 1983.
- 230 Watt, J. P., and T. J. Ahrens, Shock wave equations of state using mixed-phase regime data, *J. Geophys. Res.*, 89, 7836-7844, 1984.
- 231 Watt, J. P., and T. J. Ahrens, Shock-wave equation of state of enstatite, *J. Geophys. Res.*, 91, 7495-7503, 1986.
- 232 Yagi, T., K. Fukuoka, and Y. Syono, Shock compression of wüstite, *J. Geophys. Res. Lett.*, 15, 816-819, 1988.
- 233 Zubarev, V. N., and G. S. Telegin, The impact compressibility of liquid nitrogen and solid carbon dioxide, *Sov. Phys. Doklady*, 7, 34-36, 1962.

